

**PRELIMINARY NOTES ON THE REPRODUCTIVE BIOLOGY OF
THE LIZARDFISH, *SYNODUS SAURUS* (ACTYNOPTERYGII: SYNODONTIDAE)
IN THE AZORES**

by

Luís SOUSA (1), João Pedro BARREIROS (1), Marta S.C. SOARES (1),
Maurício HOSTIM-SILVA (2) & Ricardo S. SANTOS (3)

ABSTRACT. - Between March and November 2000, 307 specimens of the lizardfish *Synodus saurus* (Linnaeus, 1758) were captured on several coastal areas of Terceira Island, Azores. This species is an important coastal epibenthic predator. Although common, its biology, namely its reproduction, is virtually unknown. During the study period, 206 females (155 mm to 460 mm TL) and 101 males (269 mm to 290 mm TL), were captured. Sex ratio was 2:1 with more males in June. Gonadosomatic Index (GSI) for females and males, attained its maximum values during Spring and Summer, showing that these are the most intensive reproduction periods. Hepato-somatic Index (HSI) suggests that this species may well recur on body reserves for energy metabolization in gonad development. Nevertheless, more studies are necessary in order to confirm this assumption. Ovarian histology, presenting different oocytarian phases, indicates that this species has an asynchronous posture. The male increment of June, associated to higher GSI values, suggests that this could represent a moment of intense competition for females, within the studied area.

RÉSUMÉ. - Notes préliminaires sur la reproduction du poisson lézard, *Synodus saurus* (Actinopterygii : Synodontidae) aux Açores.

Entre mars et novembre 2000, 307 poissons lézard, *Synodus saurus* (Linnaeus, 1758) ont été capturés à divers endroits de la côte de l'île Terceira, aux Açores. Cette espèce est un important prédateur épibenthique. Bien qu'elle soit une espèce assez commune, sa biologie, et surtout sa reproduction, est virtuellement inconnue. Pendant cette étude, 206 femelles (de 155 mm à 460 mm LT) et 101 mâles (de 269 à 290 mm LT) ont été capturés. Le sex ratio était de 2:1, avec plus de mâles en juin. L'indice gonado-somatique (GSI) pour les femelles et les mâles, a atteint ses valeurs les plus élevées au printemps et en été, ce qui montre que ces mois constituent la période de reproduction la plus intense. L'indice hépato-somatique (HSI), indique que les réserves corporelles sont susceptibles d'être utilisées pour le développement des gonades. Cependant, cette hypothèse doit être confirmée par d'autres études. L'histologie de l'ovaire, qui présente des phases oocytaires différentes, indique que *S. saurus* a une ponte asynchrone. L'augmentation du nombre de mâles en juin, associée aux valeurs plus élevées de GSI, suggère l'existence, à cette période, d'une intense compétition pour les femelles.

Key words. - Synodontidae - *Synodus saurus* - ANE - Azores - Reproduction - sex ratio.

Teleost fishes achieved success in distinct environments for having a large array of reproductive tactics and strategies. The great adaptive capacity of teleost fishes allowed them to overcome both abiotic and biotic changes, with the possibility to reproduce successfully in changing and unstable environments (Vazzoler, 1996). Studies of fish reproduction are biologically and ecologically important (Lam, 1983; Garcia-Diaz *et al.*, 1996) and constitute a valuable tool in assessing data for fisheries management purposes (Cadima, 1984).

Reproduction in fishes is a cyclical process (Munro, 1990), that is reflected in gonadic weight gain (Vlaming *et al.*, 1992) due to maturation, increases in liver weight, where energy reserves are stored (Love, 1970; Podroschko *et al.*, 1985) together with other parts, such as muscle, and

directly from food (Vazzoler *et al.*, 1989). Several methods were developed to study fish reproduction. Quantitative and qualitative methods were used. One of the most important quantitative methods is the Gonado-somatic Index (GSI), which allows identification of the breeding season (Erickson *et al.*, 1984), together with histological observations to verify maturation of gonads (Hunter and Macewicz, 1985). The Hepato-somatic Index (HSI) (Patzner, 1980) allows to assess the role of the liver on reproduction.

Synodontidae are represented in the Azores by the following species: *Bathysaurus ferox* Günther, 1878, *Bathysaurus mollis* Günther, 1878, and *Synodus saurus* (Linnaeus, 1758) (Santos *et al.*, 1997). The lizardfish, *S. saurus* has a cylindrical shaped body, and inhabits sandy bottoms in shallow waters (Saldanha, 1997). It is a carnivore.

(1) University of the Azores, Dpt. Agricultural Sciences, 9701-851 Angra do Heroísmo, PORTUGAL. [jpedro@angra.uac.pt]

(2) University of Vale do Itajaí, Center of Technological Sciences from the Earth and Sea, R. Uruguai, C.P. 360, 88302-202 Itajaí, SC, BRAZIL.

(3) University of the Azores, Dpt. Oceanography and Fisheries, 9901-862 Horta, PORTUGAL.

rous species that eats mainly small pelagic fish, such as *Sardina pilchardus* (Soares, 2001), and occasionally, crustaceans and molluscs, using both ambush and chase tactics (Golani, 1993). Only Golani (1983) described a reproductive period between February and August in the Mediterranean. Its reproductive biology is widely unknown.

The species occurs in the Eastern Atlantic (Azores, Madeira, Cabo Verde and Canary islands), Mediterranean and Morocco Atlantic shelf waters. In the western Atlantic it is known from the Bermudas, Bahamas and Leeward islands (Sulak, 1989).

Here we give preliminary results on the reproductive biology from a sample of *S. saurus* caught in Terceira Island, Azores (NE Atlantic).

MATERIAL AND METHODS

Specimens were collected by spearfishing between March and November 2000 in Terceira island, Azores (Fig. 1). This method makes selection easy (Derbal and Kara, 1996) and was used in previous works (e.g., Barreiros and Santos, 1998)

Specimens were measured to the nearest millimeter (Total length - TL and Standard length - SL) and weighed with a precision of 0.01g. Sex determination was only possible after gonad removal. Gonads were preserved in Bouin solution for 24 to 48 h, and transferred to ethanol 70% (Vazzoler, 1996). Sections were taken from the medial part




Figure 1. - The Azores Islands (NE Atlantic) - © ImagDOP.

of the left gonad and embedded in parafin. Gonads were histologically prepared and Gonado-somatic (GSI) and Hepato-somatic (HSI) indexes were determined both for females and males. Sectioned at 2 μm, and stained with “Harris” hematoxylin and eosin, each sample was examined microscopically to determine the reproductive state. Classification of gonads followed an adaptation based on the maturity criteria of Budnichenko and Dimitrova (1981), Vazzoler (1996) and Lourinho (1998). Ovarian and testis maturation scales are presented in table I.

Gonado-somatic index (GSI), was calculated as (Vazzoler, 1996):

$$GSI = (GW/(TW-GW)) \times 100,$$

where GW is the gonad weight (g) and TW is the total fish weight (g).

Hepato-somatic index (HSI) was calculated as (Patzner, 1980):

$$HSI = (LW/TW-LW) \times 100,$$

where LW is the liver weight (g) and TW is the total fish weight (g).

Average indexes were plotted to show seasonal reproductive pattern. Sea surface temperature was measured in all sampling dates with a thermometer.

RESULTS

Two hundred and six females were captured ranging in size from 155 mm to 460 mm TL. The commonest size range was between 307 mm and 338 mm TL (SE = 3.6). Similarly, 101 males were captured during the study period. Their commonest size ranged between 269 mm and 290 mm TL (SE = 4.7), with a minimum at 185 mm to a maximal at 395 mm TL. The sex ratio observed was 2:1. In June the number of males increased turning the sex ratio from 2:1 to 1:1.

Water surface temperature gave average values somehow atypical, reaching a maximum value in October (20.7°C) (Fig. 2).

For females, GSI results showed a pronounced growth reaching a maximal peak in July (8.26). From July to August, GSI has an evident decrease, reaching a minimal value in October (0.75). HSI has an increase until May (2.06), followed by a maximal peak in July (2.26), with a

Females		Males	
Stage A	Immature	Stage A	Immature
Stage B	In Maturation	Stage B	In Maturation
Stage C	Mature	Stage C	Mature
Stage D	In recovery post-spawning	Stage D	In recovery
Stage Bp	Ovaries in maturation	Stage Bp	In maturation after ripe
Stage Cp	Mature Ovaries		

Table I. - Gonad maturation scale used for *S. saurus*. Adapted from Lourinho (1998).

minimum in August (1.65) (Fig. 3). For males, GSI has a maximal peak in June (3.96) and a minimal value in October (0.71). The HSI has several peaks during the study period. Two, less intense peaks occurred in April (1.76) and in July (1.89), and were followed by a maximal peak in September (2.52). In November, it reaches the maximal value of 2.90 (Fig. 4).

The percentage of the maturing stage for females after histological observation is presented in figure 5. Stages C and Cp represent the maximal ovarian development, and show that the percentage of mature individuals, increases until 100% in July. In October it reaches a minimum (9.5%), and increases again in November (to 38%). Testis histological observation produced the following results for each maturation stage (Fig. 6). The percentage of individuals on maturation stage C, shows that 50% of mature males occur in March, decreasing to a minimal in May with only 10%.




Figure 2. - Average temperatures of sea water during sampling period.

maximal value is reached (93%).




Figure 3. - GSI (□) and HSI (Δ) variation for females of *S. saurus* (mean values).




Figure 4. - GSI (□) and HSI (Δ) variation for males of *S. saurus* (mean values).




Figure 5. - Distribution of the maturation stages (%) for females of *S. saurus* during the collection period. A - Immature, B1 - Beginning of maturation, B2 - End of maturation, C - Mature, D - In recovery post-spawning, Bp - Ovaries in maturation, Cp - Mature ovaries.




Figure 6. - Distribution of the maturation stages (%) for males of *S. saurus* during the collection period. Abbreviations as in figure 5. Bp - Mature testicles.

DISCUSSION

GSI results for females, suggest that *Synodus saurus* in the Azores have a more intensive reproduction period during Spring and Summer, with a peak in July. This fact is confirmed by histological observation. The maximal average value obtained for females was 8, which, according to Tyler and Sumper (1996), indicates asynchronous spawning cycles. GSI for males suggest that, like females, the most intense reproduction period is during Spring and Summer. However, the histological observation did not confirm this fact and the percentage of males in stage C in this period is smaller than the values found during Autumn months. The cause of this can be due to a smaller number of specimens sampled during Spring and Summer which made the histological results inconsistent with the analytic results.

It is known that energy stored in the liver, both in males and in females, is channelled to muscular activity and for reproduction effort, being more evident in females, where the lipidic reserves are later transferred to oocytes for initial embryo development (Love, 1970). HSI results for females, during the more intensive reproduction period (July), have a peak, suggesting that the liver reserves may not be used in the final maturation stages. Nevertheless, care must be taken on this observation, because accumulation and storage of fat within the liver, in the course of ovary maturation, is known in many fish species (Bailey, 1952; Smith, 1957).

For males, the minimal value obtained in June, when GSI is highest, suggest that the liver has a weight loss, which may indicate the mobilization of hepatic reserves for testis maturation. However, this mobilization could also be derived to body condition. Once again this must be viewed with care. Other methods would be needed in order to test this hypothesis such as hepatic histological analysis.

The maturation type found in females shows similar macroscopic and microscopic characteristics to those observed by Budnichenko and Dimitrova (1981) in the con-familial species *Saurida tumbil* and *S. undosquamis*. The resemblance is in the type of development until it gets to a mature stage. In this are included all stages of oocyte development until spawning and the type of gonad coloration during that process. Budnichenko and Dimitrova (1981) described maturation stages after spawning, where it was possible to find oocytes in early and late vitellogenesis, referring to *Saurida tumbil* and *S. undosquamis* as having intermittent spawning periods. The continuous maturation of several generations of oocytes during the spawning season provides evidence of an unbroken type of maturation (Oven, 1976).

It appears that males have continuous sperm production cycles, where they ripe, and simultaneously start a new cycle of sperm production. This fact was already described in other teleost fish, such as *Ophidion* sp. (Mattei *et al.*,

1993). The percentage of mature males during Spring and Summer was too low and inconsistent with GSI results. This can be connected with the fact that the number of males collected during Spring and Summer was not enough to achieve consonance between histological observations and GSI results. The high percentage of mature males in October and November and mature females in October and September, may be explained by the higher temperatures verified in those months, probably causing an unusual maturation period, that could only be detected by histological analysis.

The role of liver reserves during the reproduction effort did not become clear.

Our results indicate that the increment of males during June, one month before the maximal GSI of males, can be due to some kind of competition strategy between males resulting in the selection of the more fit for reproduction.

This work leads to new research possibilities that are important to complement the information exposed above. It is necessary to collect more specimens for longer periods in order to test our results and detect eventual seasonal and/or inter-annual variations. The small size of our sample did not allow an efficient analysis of data from different size classes. It is also important to produce age-determination studies for this species.

The results presented in this paper should be considered as preliminary.

Acknowledgments. - We thank Alfredo Borba, for logistic support. Hernâni Martins, for the use of Histopathology equipments, and to Durval Costa, for the huge help. We thank Pedro Barreiros, and the two anonymous referees for their comments and suggestions, which greatly improved previous versions of the ms. This project was partly financed by the Departamento de Ciências Agrárias of the Universidade dos Açores.

REFERENCES

- BAILEY B.E., 1952. - Marine oils with particular reference to those of Canada. *Fish. Res. Can. Bull.*, 89: 1-413.
- BARREIROS J.P. & R.S. SANTOS, 1998. - Notes on the food habits and predatory behaviour of the dusky grouper, *Epinephelus marginatus* (Lowe, 1834) (Pisces: Serranidae) in the Azores. *Arquipélago. Life Mar. Sci.*, 16A: 29-35.
- BUDNICHENKO V.A. & O.S. DIMITROVA, 1981. - The reproductive biology of *Saurida undosquamis* and *Saurida tumbil* (Family Synodontidae) in the Arabian Sea. *J. Ichthyol.*, 19: 80-86.
- CADIMA E.L., 1984. - Avaliação do estado de um recurso e da pesca. Seminário conjunto Mocambique/NOPAD sobre "Os Recursos Marinhos de Mocambique". *Rev. Inv. Pesq.*, 9: 141-177.
- DERBAL F. & M.H. KARA, 1996. - Alimentation estivale du mérrou, *Epinephelus marginatus* (Serranidae), des côtes est Algériennes. *Cybium*, 20(3): 295-301.

- ERICKSON L.D., HIGHTOWER E.J. & D.G. GROSSMAN, 1984. - The Relative Gonadal Index: An alternative index for quantification of reproductive condition. *Comp. Biochem. Physiol.*, 78A: 151-156.
- GARCÍA-DÍAZ M.M., SANTANA J.L., TUSET V.M., RICO V. & J.A. GONZÁLEZ, 1996. - Aspectos reproductivos de *Serranus atricauda* y *S. cabrilla* (Osteichthyes, Serranidae) en Canarias. In: IX Simposio Iberico de Estudios del Bentos Marino, pp. 90-91. Resúmenes.
- GOLANI D., 1993. - The biology of the Red Sea migrant, *Saurida undosquamis* in the Mediterranean and comparison with the indigenous confamilial *Synodus saurus* (Teleostei: Synodontidae). *Hydrobiol.*, 271: 109-117.
- HUNTER J.R. & B.J. MACEWICZ, 1985. - Rates of atresia in the ovary of captive and wild northern anchovy, *Engraulis mordax*. *Fish. Bull.*, 83: 119-136.
- ISAAC-NAHUM V.J. & A.E.A. DE M. VAZZOLER, 1987. - Biología reproductiva de *Micropogonias furnieri* (Desmarest, 1823) (Teleostei, Sciaenidae). 2. Relação gonado-somática, comprimento e peso dos ovários como indicadores do período de desova. *Bol. Inst. Ocean. São Paulo*, 35(2): 123-134.
- LAM T.J., 1983. - Environmental influences on gonadal activity in fish. *Fish Physiol.*, 9B: 65-115.
- LOURINHO P.M., 1998. - Estudo da reprodução da garoupa *Serranus atricauda* (Günther, 1974). Estágio de Licenciatura em Engenharia Zootécnica, 48 p. Departamento de Ciências Agrárias. Univ. dos Açores.
- LOVE R.M., 1970. - The Chemical Biology of Fishes. 547 p. London: Academic Press.
- MATTEI X., SIAU Y., THIAW O.T. & D. THIAM, 1993. - Peculiarities in the organization of testis of *Ophidion* sp. (Pisces Teleostei). Evidence for two types of spermatogenesis in teleost fish. *J. Fish Biol.*, 43(6): 931-937.
- OVEN L.S., 1976. - Features of the Oogenesis and the Nature of Spawning of marine Fishes. 321 p. Kiev: Naukova Dumka Press.
- PATZNER R.A., 1980. - Cyclical changes in weight and fat content of the liver and their relationship to reproduction in the Hagfish *Eptatretus burgeri* (Cyclostomata). *Acta Zool.*, 61: 157-160.
- PODROSCCHKO S., PATZNER R.A. & H. ADAM, 1985. - The reproduction of *Blennius pavo* (Teleostei, Blenniidae). IV. Seasonal variation in HSI, the liver glycogen value and histological aspects of the liver. *Zool. Anz.*, 215(5/6): 256-273.
- SALDANHA L., 1997. - Fauna Submarina Atlântica (3rd edit.). 270 p. Lisboa: Publicações Europa América.
- SANTOS R.S., PORTEIRO F.M. & J.P. BARREIROS, 1997. - Marine fishes of the Azores. Annotated checklist and bibliography. *Arquipélago. Life Mar. Sci.*, Supp. 1: 1-231.
- SMITH S., 1957. - Early development and hatching. In: The Physiology of Fishes, Vol. I (Brown M.E., ed.), pp. 323-359. New York: Academic Press.
- SOARES M., 2001. - Ecologia alimentar e aspectos comportamentais de *Synodus saurus* (Linnaeus, 1758) (Actinopterygii: Synodontidae) dos Açores. Estágio de Licenciatura em Biologia Aplicada aos Recursos Animais Variantes Marinhos, 49 p. Departamento de Zoologia e Antropologia. Faculdade de Ciências da Universidade de Lisboa.
- SULAK K.J., 1989. - Synodontidae. In: Fishes of the North-eastern Atlantic and Mediterranean, Vol. I (Whitehead P.J.P., Bauchot M.-L., Hureau J.-C., Nielsen J. & E. Tortonese, eds), pp. 405-411. Paris: UNESCO.
- TYLER C.R. & J.P. SUMPTER, 1996. - Oocyte growth and development in teleosts. *Rev. Fish Biol. Fish.*, 6: 287-318.
- VAZZOLER A.E.A. DE M., 1996. - Biologia da Reprodução de Peixes Teleósteos. Teoria e Prática. 169 p. Maringá, São Paulo: EDUEM/SB.
- VAZZOLER A.E.A. DE M., CARACIOLO-MALTA M.C. & S.A. AMADIO, 1989. - Aspectos biológicos dos peixes amazônicos. XII. Indicadores quantitativos do período de desova das espécies do género *Semaprochilodus* (Characiformes, Prochilodontidae) do baixo rio Negro, Amazonas, Brasil. *Rev. Bras. Biol.*, 49(1): 175-181.
- VLAMING V., GROSSMAN G. & F. CHAPMAN, 1982. - On the use of Gonado-somatic Index. *Comp. Biochem. Physiol.*, 73A(1): 31-39.

Reçu le 6 février 2002.

Accepté pour publication le 26 septembre 2002.