

SHORT COMMUNICATION

Sponge perforating lace coral with anticancer activity

SILVIA LINO, J.R. XAVIER, R.S. SANTOS & A. COLAÇO



Lino, S., J.R. Xavier, R.S. Santos & A. Colaço 2015. Sponge perforating lace coral with anticancer activity. *Arquipélago. Life and Marine Sciences* 32.

Silvia Lino (e-mail: silvialino@uac.pt), Ricardo S. Santos & Ana Colaço, MARE – Marine and Environmental Sciences Centre, and Department of Oceanography and Fisheries, University of the Azores, PT-9901-862 Horta, Portugal; Joana R. Xavier, Centre for Geobiology and Department of Biology, University of Bergen, Thor Møhlensgate 53A/B, NO-5020, Bergen, Norway.

This short note reports results from a pilot study to investigate new anticancer agents from deep-sea corals in which colonizing sponges were encountered. The pure white stylasterid coral fans of *Errina dabneyi* (Pourtales, 1871) are a conspicuous feature on the upper bathyal slopes in Azorean waters and can be found in depths from 215 to more than 500 m (Wisshak et al. 2009; Braga-Henriques et al. 2013). From the 26 species of *Errina* known worldwide (most from deeper waters) (Cairns 1983), *E. dabneyi* seems to be endemic to the Azores Archipelago and the adjacent Mid-Atlantic Ridge region (Zibrowius & Cairns 1992; Braga-Henriques et al. 2013).

The anticancer activity of the lipid extract from the lace coral *Errina dabneyi* (Figure 1A) was being investigated as part of a broader ongoing bio prospecting research on deep-sea invertebrates from the Azores area. Deep-sea fisheries in the Azores use longline gears at depths where cold-water corals are common (Braga-Henriques et al. 2011; Pham et al. 2014). The specimens studied were collected accidentally (bycatch) at Princess Alice Seamount (near the Azorean central group of islands) at 200 m in April 2012 by a local commercial fishery fleet. Observers onboard froze samples immediately after collection and kept them at -20°C until being extracted. A subsample was taken and preserved for the biological reference collection at University of the Azores (DOP/UAc COLETA), with ID number - DOP

9044. When processing *Errina dabneyi*, we noted that part of the coral (from the base to the middle of the primary branches) had been bored by what appeared to be a sponge in the interior of the skeleton (Figure 1B). Since it was impossible to fully separate the sponge from the coral, the part of the coral that was colonized was treated as an independent sample (CS; Figure 1c). For identification purposes, a small fragment of sponge tissue was excised, digested in sodium hypochlorite, and washed in a water-ethanol series. The obtained spicules were then mounted in Canada balsam, observed and measured under a microscope. The sponge was identified as *Thoosa armata* Topsent, 1888 (Demospongiae, Astrophorida), a species previously known to occur associated with deep-sea corals in the Azores (Topsent 1904). *Thoosa* is one of several demosponge genera known for their ability to excavate calcareous substrates (see review in Cairns 1983), a genus of demosponge in the family Thoosidae. This genus is known for boring holes in corals (Schoenberg 2008) as can be seen in Figure 1B. Total lipids were extracted using a modified Bligh and Dyer method as described in Lino et al. (2013). Cytotoxicity essays were performed on HCT-116 human colon adenocarcinoma (tumorous) cell line. Briefly, cells were inoculated into plates at 2.5×10^4 cells/ml (150 μ l/well), and incubated overnight 37°C and 5% CO₂ before treatment. Series of dilutions in DMSO of each of the




Fig. 1. a) *Errina dabneyi* specimen deposited in the reference collection at the University of the Azores DOP/UAg COLETA); b) *Errina dabneyi* specimen where it is possible to see the sponge inside (pointed by the black arrow); c) Specimen of coral *E. dabneyi* plus sponge (CS) investigated for their anticancer potential in this study - the white arrows indicate the holes done by the *Thoosa* sponge in the coral skeleton in order to connect with the exterior and which indicates the presence of the sponge within the coral.

extracts (*Errina dabneyi* (C) and *Errina* plus sponge (CS)), were added to the cells followed by further incubation at 37°C for 72 h. Each extract was assayed in duplicate. Cell viability was determined by a colorimetric [3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphen-yl)-2-(4-sulfophenyl)-2H-tetrazolium] (MTS) assay. After the exposure time, 20 µL of MTS/ PMS (phenazine methosulfate - electron coupling agent) solution was added to each well. Plates were incubat-

ed for 3 h before absorbance was measured at 490 nm using an EMax® Endpoint ELISA Plate Reader. Cells in DMSO were used as negative control, and cells treated with Etoposide (VP-16) were used as positive control. Relative cellular survival was determined by using the measured optical density (OD) and was calculated as follows: (% Survival for sample) = [(OD of sample)/(Average of negative control)] x100. Concentration that killed 50% of the cancer cells

(IC₅₀ values) was automatically calculated by SoftMax Pro software. Results for anticancer activity showed that the extract from *Errina dabneyi* did not show any activity but the extract from the coral sample harbouring the sponge was active presenting IC₅₀ values of 25.29 and 32.72 µg/mL for the two essays. To our knowledge this is the first time that anticancer activity is reported in a coral perforating sponge.

Considering the interesting preliminary results on the anticancer activity for the sponge and taking in consideration the high bio eroding ratio observed in *Errina dabneyi* specimens deposited in the collection at University of the Azores, more consistent studies should be made to address this matter.

ACKNOWLEDGMENTS

The authors would like to thank the observer onboard Daniel Pereira for bringing the samples in good preservation conditions. We are grateful to Professor William Fenical at Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography, University of California San Diego, who kindly accepted Silvia Lino as collaborator at his lab to investigate the natural marine products of our samples and to Lauren Paul for performing the cytotoxicity essays. We thank Valentina Matos and Andreia Braga-Henriques for assisting with sampling and coral taxonomic identification. Silvia Lino was supported by PhD grant from Fundação para a Ciência e Tecnologia, (FCT - Ministry of Science and Technology of Portugal), with reference SFRH/BD/72154/2010.

REFERENCES

- Braga-Henriques, A., M. Carreiro-Silva, F. Porteiro, V. Matos, Í. Sampaio, O. Ocaña, & S.P. Ávila 2011. The association between a deep-sea gastropod *Pedicularia sicula* (Caenogastropoda: Pediculariidae) and its coral host *Errina dabneyi* (Hydrozoa; Stylasteridae) in the Azores. *ICES Journal of Marine Sciences* 68: 399–407.
- Braga-Henriques, A., F. M. Porteiro, P.A. Ribeiro, V. de Matos, I. Sampaio, O. Ocaña, & R.S. Santos 2013. Diversity, distribution and spatial structure of the cold-water coral fauna of the Azores (NE Atlantic). *Biogeosciences* 10: 4009–4036.
- Cairns, S.D. 1983. A generic revision of the *Stylasterina* (Coelenterata: Hydrozoa). Part 1. Description of the genera. *Bulletin of Marine Science* 33: 427–508.
- Lino, S., B. Solé, V. de Matos, R. Pereira, J. Xavier, R. Santos & A. Colaço 2013. Lipid extractions from marine deep-sea invertebrates: modifications to the Bligh & Dyer method to increase lipid yields. *Arquipelago. Life and Marine Sciences* 31: 9–14.
- Pham C.K., H. Diogo, G. Menezes, F. Porteiro, A. Braga-Henriques, F. Vandepierre & T. Morato. 2014. Regulated bottom longline helps achieving sustainability of deep-sea fisheries. *Scientific Reports*. 4:4837.
- Schoenberg, C.H.L. 2008. A history of sponge erosion: from past myths and hypotheses to recent approaches. Pp. 165–202 in: Wishak, M. & L. Tapanila (Eds). *Current developments in bioerosion*. Erlangen Earth Conference Series. Berlin, Heidelberg: Springer-Verlag.
- Topsent, E. 1904. Spongiaires des Açores. *Résultats des campagnes scientifiques accomplies par le Prince Albert I. Monaco* 25: 1–280, pls 1–18.
- Wisshak, M., M. López Correa, H. Zibrowius, J. Jakobsen & A. Freiwald 2009. Skeletal reorganisation affects geochemical signals, exemplified in the stylasterid hydrocoral *Errina dabneyi* (Azores Archipelago). *Marine Ecology Progress Series* 397: 197–208.
- Zibrowius, H. & S.D. Cairns 1992. Revision of the northeast Atlantic and Mediterranean Stylasteridae (Cnidaria: Hydrozoa). *Mémoires du muséum national d'histoire naturelle A* 153:1–136.

Received 12 Dez 2014. Accepted 06 Feb 2015,
Published online 13 February 2015.