



UNIVERSITY OF THE AZORES

DEPARTMENT OF OCEANOGRAPHY AND FISHERIES

Effects of natural mortality on the yield per recruit models

A thesis presented to the Department of Oceanography and Fisheries at the University of the Azores in partial fulfilment of the requirements for the degree in Master of Integrated Studies of the Oceans

Ana Maria Novoa Pabon

HORTA

2015

UNIVERSITY OF THE AZORES

DEPARTMENT OF OCEANOGRAPHY AND FISHERIES

Effects of natural mortality on the yield per recruit models

**A thesis presented to the Department of Oceanography and Fisheries at the
University of the Azores in partial fulfilment of the requirements for the degree
Master in Integrated Studies of the Oceans**

Ana Maria Novoa Pabon

ADVISOR:

Mário Rui Pinho

CO-ADVISOR:

Pedro Conte de Barros

HORTA

2015

This work resulted in the following publications:

Novoa-Pabon, A., Barros, P., Krug, H & M. Pinho. 2015. Yield per recruit analysis for the black spot seabream (*Pagellus bogaraveo*) stock from the Azores (ICES area Xa2). ICES Working Document WGDEEP 2015. 24p.

Pinho, M., **Novoa-Pabon, A.**, Pereira, J. & H. Krug. 2015. Catch curve analysis for the red blackspot seabream (*Pagellus bogaraveo*) stock from the Azores (ICES Xa2). ICES Working Document WGDEEP 2015. 19p.

Pinho, M., **A. Pabon** & H. Krug. 2015. Natural mortality for Thornback (*Raja clavata*) stock from the Azores (ICES Xa2). ICES Working Document WGEF 2015. 15p.

Novoa-Pabon, A., Barros, P., Krug, H., Silva, P. & M. Pinho. 2015. Yield per recruit analysis for the red blackspot seabream (*Pagellus bogaraveo*) stock from the Azores (ICES area Xa2). ICES Annual Science Conference, ID 277 A: 33, 21th to 25th September, 2015, Copenhagen, Denmark. <http://www.ices.dk/sites/pub/ASCExtendedAbstracts/Shared%20Documents/A%20-%20Advancement%20of%20stock%20assessment%20methods%E2%80%8B%20for%20sustainable%20fisheries%E2%80%8B%E2%80%8B/A3315.pdf>

Silva, P., Krug, H., **Novoa-Pabon, A.** & Pinho, M.R., 2015. Natural mortality for the red blackspot seabream (*Pagellus bogaraveo*) stock from the Azores (ICES Xa2). ICES Annual Science Conference, ID 171 A: 28, 21th to 25th September, 2015, Copenhagen, Denmark. <http://www.ices.dk/sites/pub/ASCExtendedAbstracts/Shared%20Documents/A%20-%20Advancement%20of%20stock%20assessment%20methods%E2%80%8B%20for%20sustainable%20fisheries%E2%80%8B%E2%80%8B/A2815.pdf>

Pinho, M., **Novoa-Pabon, A.**, Pereira, J., Silva, P. & H. Krug. 2015. Catch curve analysis for the red blackspot seabream (*Pagellus bogaraveo*) stock from the Azores (ICES Xa2). ICES Annual Science Conference, ID 168 A: 27, 21th to 25th September, 2015, Copenhagen, Denmark. <http://www.ices.dk/sites/pub/ASCExtendedAbstracts/Shared%20Documents/A%20-%20Advancement%20of%20stock%20assessment%20methods%E2%80%8B%20for%20sustainable%20fisheries%E2%80%8B%E2%80%8B/A2715.pdf>

Pinho, M., **Novoa-Pabon, A.**, Silva, P., Krug, H & J. Pereira. 2015. Mixed demersal/deep-water hook and line small scale fishery from the Azores (ICES Xa2). ICES Annual Science Conference, ID 305 F: 30, 21th to 25th September, 2015, Copenhagen, Denmark. <http://www.ices.dk/sites/pub/ASCExtendedAbstracts/Shared%20Documents/F%20-%20Small-scale%20fisheries%20under%20data-limited%20scenarios/F3015.pdf>

Abstract

Natural mortality (M) is one of the most important parameter used on the Yield Per Recruit models and is commonly assumed as constant over time. However, M is strongly related to age, length or weight. This study analyses for the first time the effects of natural mortality in the yield per recruit estimates for the red black spot seabream (*Pagellus bogaraveo*) stock from the Azores archipelago. Input data were reviewed and length base YPR model was adopted in order to analyse the effect of M values, as input parameters, for constant M for all lengths (M_C) and size-specific natural mortality (M_L). Results show that estimated M_L for the *P. bogaraveo* decreases exponentially, with high mortality for juveniles ($M=0.6$ for recruits) and low mortality for the adults ($M\approx 0.2$ for Fork length >30 cm). YPR curve at long term is virtually asymptotic for any option of M which the maximum is not possible to attain except for low level of $M=0.2$. Fishing mortality corresponding to maximum sustainable yield (F_{max}) is not considered a useful target reference point for this species, because spawning biomass is usually very low (Spawning Potential Ratio - SSBPR $<20\%$) at this level of fishing mortality. In general, increasing M_C produces a clear decrease on the cohort values by length for all stock characteristics (N/R, B/R, SSB/R, C/R and Y/R). For M_L hypotheses similar trends are observed on the behaviour of the stock characteristics but with important reductions on the maximum values. At the $F_{0.1}$ strategy the yield value is lower as M increases but greater fishing mortality is required to produce the correspondent yield ($Y_{0.1}/R$). Similar to the catch higher M implies higher depletion of the abundance (B or SSB). YPR analysis indicates that *P. bogaraveo* from the Azores is overexploited ($F_{current} 53\% > F_{0.1}$ and SSBPR $<20\%$). Changing the natural mortality does not change the perception of the stock status, with M_L implying higher reductions on the stock characteristics. The distribution of M_L has an effect similar to increase the M_C . However, the problem is how to reconcile the M_L with M_C . Under the same relative exploitation pattern (L_c), increasing M_C has a high effect on YPR, reducing the stock characteristics and increasing the fishing mortality required to achieve the maximum sustainable catch. Implications of these variations for the stock assessment are discussed.

Keywords: Stock assessment, Yield per recruit, natural mortality, *Pagellus bogaraveo*, Azores.

Resumo

A mortalidade natural (M) é um dos mais importantes parâmetros utilizado nos modelos de captura por recruta e é geralmente assumido como constante ao longo do tempo. No entanto, M é fortemente relacionada com a idade, comprimento ou peso. Este estudo analisa pela primeira vez os efeitos da mortalidade natural na estimativa da captura por recruta para o stock de Goraz (*Pagellus bogaraveo*) no Arquipélago dos Açores. Os dados de entrada foram revistos e o modelo de YPR baseado no comprimento foi adotado a fim de analisar o efeito de M como parâmetros de entrada, considerando M constante para todos os comprimentos (M_C) e mortalidade natural específico por comprimentos (M_L). Os resultados mostram que M_L estimado para *P. bogaraveo* diminui exponencialmente, com elevada taxa de mortalidade nos jovens ($M = 0,6$ para recrutas) e baixa mortalidade para os adultos ($M \approx 0,2$ para comprimento furcal $FL > 30\text{cm}$). A curva de YPR a longo prazo é praticamente assintótica para qualquer opção de M , o máximo não é possível atingir com exceção para baixos valores de $M = 0,2$. A mortalidade por pesca referente á captura máxima sustentável (F_{\max}) não é considerado um ponto de referência alvo útil para a espécie, porque a biomassa desovante (proporção potencial de desova) correspondente é normalmente muito baixa ($SSBPR < 20\%$). Em geral, o aumento de M_C produz uma clara diminuição sobre os valores da coorte por comprimento para todas as características do stock. Para a hipóteses M_L tendências semelhantes são observados no comportamento das características do stock (N/R , B/R , SSB/R , C/R e Y/R), mas com reduções importantes em relação aos valores máximos. Na estratégia $F_{0.1}$ o valor de captura é inferior com o aumento de M , mas maior mortalidade por pesca é necessária para produzir a captura correspondente ($Y_{0.1}/R$). Semelhante à maior captura M implica maior esgotamento da abundância (B ou SSB). A análise de YPR para *P. bogaraveo* dos Açores indica que o stock está sobreexplorado ($F_{\text{actual}} 53\% > F_{0.1}$ e $SSBPR < 20\%$). Alterar a mortalidade natural não muda a percepção do estado do stock e a adopção de M_L a implica reduções mais elevadas sobre as características do stock. A distribuição de M_L tem um efeito semelhante ao de aumentar a M_C . No entanto, o problema é como conciliar a M_L com M_C . Sob o mesmo padrão relativo de exploração (L_c), o aumento do M_C tem um alto efeito sobre YPR, reduzindo as características do stock e o aumento da mortalidade por pesca necessária para alcançar a captura máxima sustentável. São discutidas implicações destas variações para a avaliação do recurso.

Palavras-chave: Avaliação de recursos, captura por recruta, mortalidade natural, *Pagellus bogaraveo*, Açores

Acknowledgments

This thesis is the result of a year of illusions, satisfactions, work and patience, and it would not have been possible without the help of many people.

First of all, I would like to thank my supervisors for their guidance and support to delve every day into the world of fishing. In particular, I would like to thank Dr. Mario Pinho for his optimism and especially for his patience with all my questions, and Dr. Pedro Barros for his questions that always made me think a little deeper about my work.

I am also grateful to the Department of Oceanography and Fisheries at the University of the Azores (DOP/UAç) for the opportunity to be part of this institution, and by making me feel part of this family. I am also gratified by being allowed to stay in the Laboratory of Oceanography where I always felt at ease and welcome. I thank Prof. Ana Martins for her help and understanding. I would like to extend my thanks to Rui Rosa for allowing me to join me in his work and for his enthusiasm to teach me, as well as, for the early mornings in the auction. Priscilla, I could not have had a better friend and colleague and as you say: "Friends are everything."

I am most grateful to my family for always helping and supporting me in every step of the way. In particular, I would like to thank my parents to whom I owe everything, especially for their love and constant support throughout this time, and by never letting me give up while always pushing me to fulfil my dreams. Dear sister, I am also very thankful for making me always feel that the great distance among us was actually much shorter.

To my friends who are both close and far away, thank you so much all for being always there for me during both good and bad times. You have taught me that the family can be as large as you want it to be.

I would also like to acknowledge the staff at DOP/UAç for all the administrative and logistics help along the way. This is greatly appreciated.

Last but not least, I would sincerely like to leave a grateful note to the jury members present in my thesis defense and for the time they have taken to read and listen to my work.

To all, I do thank you very much for your patience and care, since you know that, without you I would have never made it.

Index

Abstract	IV
Resumo.....	V
Acknowledgments.....	VI
Index.....	VII
Figure Index.....	IX
Table Index	X
Introduction.....	1
Background.....	3
1. Natural mortality (M)	3
2. Selectivity effects.....	4
3. Yield per recruit (YPR).....	5
4. Exploitation pattern.....	6
5. <i>Pagellus bogaraveo</i> (Brünnich, 1768) – Red Black Spot Seabream	7
Methods	9
1. Definition of input parameters.....	10
I. Life history.....	10
II. Natural mortality	10
III. Natural mortality constant - M_C	10
IV. Natural mortality dependent by length - M_L	10
V. Biological data.....	11
VI. Current fishing mortality.....	12
VII. Selectivity	12
VIII. Mean price per Kg	12
2. Effects of different natural mortality values on the cohort characteristics by length (N_L , B_L , SSB_L , C_L , Y_L)	12
3. Yield per recruit (YPR).....	13
4. Optimal exploitation pattern.....	13
Results	15

1. Life history	15
2. Natural mortality dependent by length - M_L	16
3. Effects of different natural mortality values on the cohort characteristics by length (N_L , B_L , SSB_L , C_L , Y_L)	17
4. Yield per recruit (YPR).....	19
I. Base case (current situation)	19
II. Effects of natural mortality constant for all length on the YPR stock characteristics	20
III. Natural mortality variable by length	23
5. Exploitation pattern.....	23
Discussion.....	27
Reference	31
Annex.....	39

Figure Index

Figure 1. Theory forms for the selection curves of the fishing gears.	5
Figure 2. Graph of selectivity model (logistic curve) (s_i), natural mortality constant for all lengths (M_C) and natural mortality variable by length (M_L).....	9
Figure 3. Natural mortality estimates of red blackspot seabream (<i>Pagellus bogaraveo</i>) from the Azores (ICES area X), using indirect methods applied to base case life history parameters of Table 2. Dashed line represents the mean value of the estimates. Shading area represents 95% confidence interval.	16
Figure 4. Behaviour of stock characteristics by length (N_L , B_L , SSB_L , Y_L and C_L) for different values of M_C and for M_L	18
Figure 5. Yield per recruit analyse for the base case (M_C). Horizontal dashed grey line represents the 20-40% Spawning Potential Ratio range.....	20
Figure 6. Effects of natural mortality independent of size (M_C) and dependent of size (M_L) in the results of yield per recruit analyse (YPR). The graph at the right bottom (U) permits to map the $F_{0.1}$ (maximum value) for each level of natural mortality. 21	
Figure 7. Yield per recruit curves for optimal length of first capture, for the two assumptions: M_C and M_L	24
Figure 8. Yield per recruit, Biomass per recruit (B/R) and Spawning biomass per recruit (SSB/R) isopleths for the assumption of M_C and M_L . Dots indicate $F_{0.1}$ values estimated for each L_c and the star indicate the $F_{current}$ for $L_c= 30\text{cm}$	26

Table Index

Table 1. Length specific YPR input parameters for <i>Pagellus bogaraveo</i> of the Azores (ICES area X). Selectivity (s_i) of the gear (Sousa <i>et al.</i> , 1999); females sex-ratio for the period 1982-1991 (% Females) from ICES (2012) ; Maturation females for the year 1991 from Krug (1998) as computed by ICES (2012); Price is for Kg in Euro.	11
Table 2. Life history parameter for <i>Pagellus bogaraveo</i> of the Azores (ICES area X2a), adopted as input parameters (base case) for YPR.	15
Table 3. Natural mortality by length, using the empirical formula of Gislason <i>et al.</i> (2010), estimated for <i>Pagellus bogaraveo</i> of the Azores. Parameters used are defined on Table 2. Life history parameter for <i>Pagellus bogaraveo</i> of the Azores (ICES area X2a), adopted as input parameters (base case) for YPR.	17
Table 4. Results of yield per recruit analysis for different management strategies and values of natural mortality independent of size (M_c) and dependent of size (M_L). Shading column correspond to the base case. N.C. - Not converge.	22

Introduction

The yield per recruit analysis (YPR) is the most commonly model used for fish stock assessment since its initiation with Beverton and Holt in 1957 (Xiao, 2002). Yield per recruit model assume that the population is in steady state conditions without major changes in mortality, growth and recruitment in the long term, but is independent of stock size (Kirchner, 2001; Xiao, 2002; Law *et al.*, 2012). Natural mortality (M) is the main parameter using in all mathematical models of fish stock dynamics and one of the most important parameter used on the YPR models. Usually it is difficult to measure natural mortality directly and for this reason is assumed in stock assessment as constant and independent for all age (Beverton & Holt, 1957; Gulland, 1965; Cadima, 2003). However, in many fish stock M is far from to be constant and this problem is extensive enough for not consider this variation (Vetter, 1988). M in general is extremely high for egg and larval stages, decrease in the juvenile, is low for old and mature fish, and increases again with senescence (Giménez-Hurtado *et al.*, 2008; Law *et al.*, 2012), indicating the variability of M with the age-structure. The main characteristic of M is to have influence on estimates of maximum yield (Y/RMAX; FMAX) as in Beverton and Holt yield models and cohort analyses (Vetter, 1988; Beyer *et al.*, 1999). However, these effects of M on the YPR models are not well addressed in the literature, particularly when related with the definition of the optimal exploitation pattern for the same M values considered as constant or when distributed by length (or ages).

The red black spot seabream (*Pagellus bogaraveo*) is considered a dominant species of the Azorean deep-water fishery (Pinho & Menezes, 2009). In the Eastern Atlantic red seabream distributed from south of Norway to Cape Blanc in Mauritania, Azores, Madeira and Canary Islands and Mediterranean Sea and in summer extends to west of Scotland (Pinho, 2003; Herrera, 2010; Lorange, 2010). Blackspot sea bream has been exploited in the Azores (Area Xa2), at least since the XVI century as part of the demersal fishery. The directed fishery is a hook and line fishery where two components of the fleet can be defined: the artisanal (hand lines) and the longliners (Pinho, 2003; Pinho & Menezes, 2005; Pinho & Menezes, 2009; Pinho *et al.*, 2014).

This stock has no management objectives defined because the uncertainty on the assessment, being precautionary managed based on trends of abundance indices and landings (ICES, 2014). Yield per recruit (YPR) analysis may be a useful tool to explore long term management objectives for red seabream and assess the actual state of the stock. Application of the YPR method to this species was already reported by Lorange (2010) for the Biscay Golf (ICES area VIII) and Pinho *et al.* (2014) for the Azores.

This study investigates for first time the effects of natural mortality in the yield per recruit estimates for the red seabream stock from the Azores archipelago. Main objectives are: a) Review the estimation procedures for natural mortality in order to evaluate the variability within and the difference between constant and size-specific M hypothesis; b) Analyse the effect of M on the behaviour of different YPR stock characteristics; c) Analyse the effect of changing M on the assessment results of Red blackspot seabream (*Pagellus bogaraveo*) stock of the Azores in order to define long term management objectives.

The following section presents a general background information about approaches for the natural mortality, gear selectivity, yield per recruit and optimal exploitation pattern often used in fisheries stock assessment models and implications in the management including a brief review of the Azorean red seabream stock and fisheries. Section 2 describes the methods undertaken, section 3 describe the results found and section 4 focus on the discussion of the results of this study.