

AGE AND GROWTH OF *TRACHURUS PICTURATUS* (BOWDICH, 1825) (TELEOSTEI: CARANGIDAE) FROM THE AZORES

HELENA AZEVEDO ISIDRO

ARQUIPÉLAGO



ISIDRO, HELENA AZEVEDO 1990. Age and growth of *Trachurus picturatus* (Bowdich, 1825) (Teleostei: Carangidae) from the Azores. - *Arquipélago*. Life and Earth Sciences 8:45-54. Angra do Heroísmo. ISSN 0870-6581.

From late 1983 to early 1987, 2530 *Trachurus picturatus* were sampled from captures around Faial and Pico Islands (belonging to the Central Group of the Azorean Archipelago). Biological parameters were taken using either fresh or frozen fish. Students' *t* test was applied to compare the slopes of the obtained weight-length regressions, leading to the conclusion that the two samples were significantly different for a 95% confidence limit. Thus parameters obtained with frozen fish were discarded. Weight-length regression obtained was $W_t(\text{g}) = 0.00819 \times L_f^{3.11}$ (cm). Total length-fork length regression was $L_t(\text{cm}) = -0.447 + 1.137 L_f$ (cm). The regressions between length, height and weight of left and right sagitta otoliths and the respective fork length were calculated. Further comparison between the obtained parameters lead to the conclusion that either could be used. Otolith sections were found to be the more reliable structures for age determination. An age-length key was established. The best fit to von Bertalanffy growth equation, using otolith age determination, was: $L_\infty = 52.9$ cm; $k = 0.20 \text{ year}^{-1}$; $t_0 = -0.23$ year. Nevertheless a Gompertz growth curve may give a better fit.

ISIDRO, HELENA AZEVEDO 1990. Idade e crescimento de *Trachurus picturatus* (Bowdich, 1825) (Teleostei: Carangidae) dos Açores. - *Arquipélago*. Ciências da Natureza 8:45-54. Angra do Heroísmo. ISSN 0870-6581.

Desde fins de 1983 até 1987 efectuou-se a amostragem biológica de 2530 *Trachurus picturatus* provenientes de capturas efectuadas perto das Ilhas do Faial e Pico (ambas pertencentes ao Grupo Central do Arquipélago dos Açores). Os parâmetros biológicos foram obtidos em exemplares frescos e congelados. O teste *t* foi aplicado para comparar os declives das duas regressões comprimento-peso obtidas, permitindo concluir que as duas amostras eram significativamente diferentes para um limite de confiança de 95%. Daí que no presente trabalho os parâmetros obtidos com exemplares congelados tenham sido omitidos. A relação comprimento-peso obtida foi: $W_t(\text{g}) = 0.00819 \times L_f^{3.11}$ (cm). A relação comprimento total-comprimento furcal foi: $L_t(\text{cm}) = -0.447 + 1.137 L_f$ (cm). Também foram calculadas as relações entre o comprimento, a altura e o peso dos otólitos (*sagittae*) esquerdo e direito, e o comprimento furcal do respectivo indivíduo. A comparação das relações obtidas levou-nos a concluir que poderíamos utilizar quer o otólito esquerdo quer o direito. Para a determinação da idade concluiu-se que as secções dos otólitos eram as estruturas mais fáceis de observar. Foi estabelecida uma chave comprimento-idade. A melhor estimativa para a equação de crescimento de von Bertalanffy, determinando a idade através de otólitos foi: $L_\infty = 52.9$ cm; $k = 0.20 \text{ ano}^{-1}$; $t_0 = -0.23$ ano. Contudo a equação de crescimento de Gompertz talvez se ajuste melhor.

Helena Azevedo Isidro, Departamento de Oceanografia e Pescas, Universidade dos Açores, P-9900 Horta, Açores, Portugal.

INTRODUCTION

Azorean fishery

Trachurus picturatus (Bowdich, 1825) (Portuguese common names are "carapau negro", in mainland and "chicharro", in Azores and Madeira) is the most important of the small pelagic species caught in the Azores. Data available since 1969 show a variation of commercial landings of horse mackerel between 2 and 4 thousand metric tons a year. From 1980 to 1987 they represented 84,5 % of the small pelagics total landings (Figure 1), mostly captured in the Oriental Group of the Azorean Archipelago.

For the same period *T. picturatus* annual landings have been in the first place in total landings (metric tons) and in first or second places in commercial value (thousand escudos) within all marine commercial Azorean species (with exception of tunas).

Besides landing data one must also consider

the captures, not quantified, as live bait for tuna fishery, highly important for the Archipelago fishery industry.

Small immature individuals of *T. picturatus* are mostly captured near the islands coasts, with dipnet or purse seine, either by the artisanal fleet or by tuna boats, respectively. Larger fishes, over 2 or 3 years old, represent an accidental capture in a longline fishery directed towards blackspot seabream (*Pagellus bogaraveo*) and other demersal species, in deeper waters and near "fishing banks".

Taxonomic position and distribution

There is no general agreement about the taxonomic position of Azorean horse mackerel.

Ocean scad (RYTOV & RAZUMOVSKAYA 1984), ocean mackerel or blue scad (HUREAU & MONOD 1979), and blue jack mackerel (SMITH-VANIZ 1986) are some common English




Fig. 1 - Weight percentage of small pelagics from commercial landings (Azores, 1980-87).

names for the species *Trachurus picturatus* (Bowdich, 1825), widely distributed along the Pacific coast of North and South America, in the Mediterranean and the Atlantic Ocean (Canaries, Azores and Madeira Archipelagos) (RYTOV & RAZUMOVSKAYA 1984) as well as in China (LETACONNOUX 1951).

ALEYEV (1957, cited in SHABONEYEV 1973) subdivides the genus *Trachurus* into several groups, according to morphological features, which are characteristic of the suitability of the individual forms of the genus *Trachurus* for a pelagic habitat. Thus, apart from *T. declivis*, *T. symmetricus*, *T. murphy* and *T. binghami*, he considers that the most oceanic and more complex from a systematic point of view is the species *T. picturatus*, and namely the sub-species *picturatus*.

SHABONEYEV, together with RYAZANTSEVA (1977) and with KOTLYAR (1979), refers that the oceanic horse mackerel, *Trachurus picturatus picturatus*, occurs in the Eastern Atlantic, along the coastal area of Western Europe, from the Bay of Biscay to the Azores and along the western coast of Africa to Mauritania, a typical form inhabiting the eastern part of the Central Atlantic. *Trachurus picturatus picturatus* is widely distributed not only on the continental shelf, but also far beyond its boundaries, in the regions of underwater elevations and seamounts, near the Canary Islands and the Azores (SHABONEYEV & KOTLYAR 1979).

Apart from the cited papers other authors may have used the name *Trachurus trachurus* when studying *T. picturatus*, which is unfortunate, as the biology of the two species might differ in many aspects (Helen Martins, personal communication). Thus one could possibly explain the lack of information about *T. picturatus*.

MATERIAL AND METHODS

This species has been randomly sampled from commercial landings, especially in the Central Group of the Archipelago since mid 1979. The present paper refers to biological data collected

from specimens captured around Faial and Pico Islands (Central Group), between late 1983 and early 1987.

Biological sampling was made with either fresh ($n = 2073$) or frozen ($n = 457$) fish. Weights were taken to the nearest 0.1 g and length measurements to the centimeter below, corresponding to the length-class.

Growth relations

Weight-length relationships W_t (g) = $a \times L_f^b$ (cm) were determined using total weight (W_t) against fork length (L_f), as many specimens presented damaged caudal fins. The parameters have been determined using a computer statistical program TODOS (Eduardo Isidro, personal communication).

The linear relationship between total length (L_t) and fork length (L_f), L_t (cm) = $a + b \times L_f$ (cm), was determined for the available data.

Eventual differences between sexes have not been tested, based on the statement of SHABONEYEV & RYAZANTSEVA (1977) that sexual dimorphism does not exist in oceanic horse mackerel. Subsequently it was assumed that all data could be used without separating the sexes.

Fresh and frozen samples

Freezing seems to have provoked an increase in weight and a decrease in length. Therefore, the slopes of the two logarithmic weight-length regressions were compared using the t test, as described by ZAR (1974), a simple method for testing the hypothesis about equality of two regression coefficients.

The statistic test is:

$$t = (b_1 - b_2) / S(b_1 - b_2)$$

$$\text{with } b = \sum xy / \sum x^2$$

where

$$S(b_1 - b_2) = \sqrt{(S^2_{y,x})_p / (\sum x^2)_1 + (S^2_{y,x})_p / (\sum x^2)_2}$$

and the pooled residual mean square is calculated as:

$$(S^2_{y,x})_p = [(\text{residual SS})_1 + (\text{residual SS})_2] / [(\text{residual DF})_1 + (\text{residual DF})_2]$$

$$\text{being residual SS} = \sum y^2 - (\sum xy)^2 / \sum x^2$$

names for the species *Trachurus picturatus* (Bowdich, 1825), widely distributed along the Pacific coast of North and South America, in the Mediterranean and the Atlantic Ocean (Canaries, Azores and Madeira Archipelagos) (RYTOV & RAZUMOVSKAYA 1984) as well as in China (LETACONNOUX 1951).

ALEYEV (1957, cited in SHABONEYEV 1973) subdivides the genus *Trachurus* into several groups, according to morphological features, which are characteristic of the suitability of the individual forms of the genus *Trachurus* for a pelagic habitat. Thus, apart from *T. declivis*, *T. symmetricus*, *T. murphy* and *T. binghami*, he considers that the most oceanic and more complex from a systematic point of view is the species *T. picturatus*, and namely the sub-species *picturatus*.

SHABONEYEV, together with RYAZANTSEVA (1977) and with KOTLYAR (1979), refers that the oceanic horse mackerel, *Trachurus picturatus picturatus*, occurs in the Eastern Atlantic, along the coastal area of Western Europe, from the Bay of Biscay to the Azores and along the western coast of Africa to Mauritania, a typical form inhabiting the eastern part of the Central Atlantic. *Trachurus picturatus picturatus* is widely distributed not only on the continental shelf, but also far beyond its boundaries, in the regions of underwater elevations and seamounts, near the Canary Islands and the Azores (SHABONEYEV & KOTLYAR 1979).

Apart from the cited papers other authors may have used the name *Trachurus trachurus* when studying *T. picturatus*, which is unfortunate, as the biology of the two species might differ in many aspects (Helen Martins, personal communication). Thus one could possibly explain the lack of information about *T. picturatus*.

MATERIAL AND METHODS

This species has been randomly sampled from commercial landings, especially in the Central Group of the Archipelago since mid 1979. The present paper refers to biological data collected

from specimens captured around Faial and Pico Islands (Central Group), between late 1983 and early 1987.

Biological sampling was made with either fresh (n = 2073) or frozen (n = 457) fish. Weights were taken to the nearest 0.1 g and length measurements to the centimeter below, corresponding to the length-class.

Growth relations

Weight-length relationships W_t (g) = $a \times L_f^b$ (cm) were determined using total weight (W_t) against fork length (L_f), as many specimens presented damaged caudal fins. The parameters have been determined using a computer statistical program TODOS (Eduardo Isidro, personal communication).

The linear relationship between total length (L_t) and fork length (L_f), L_t (cm) = $a + b \times L_f$ (cm), was determined for the available data.

Eventual differences between sexes have not been tested, based on the statement of SHABONEYEV & RYAZANTSEVA (1977) that sexual dimorphism does not exist in oceanic horse mackerel. Subsequently it was assumed that all data could be used without separating the sexes.

Fresh and frozen samples

Freezing seems to have provoked an increase in weight and a decrease in length. Therefore, the slopes of the two logarithmic weight-length regressions were compared using the t test, as described by ZAR (1974), a simple method for testing the hypothesis about equality of two regression coefficients.

The statistic test is:

$$t = (b_1 - b_2) / S(b_1 - b_2)$$

$$\text{with } b = \sum xy / \sum x^2$$

where

$$S(b_1 - b_2) = \sqrt{(S^2_{y,x})_p / (\sum x^2)_1 + (S^2_{y,x})_p / (\sum x^2)_2}$$

and the pooled residual mean square is calculated as:

$$(S^2_{y,x})_p = [(\text{residual SS})_1 + (\text{residual SS})_2] / [(\text{residual DF})_1 + (\text{residual DF})_2]$$

$$\text{being residual SS} = \sum y^2 - (\sum xy)^2 / \sum x^2$$




Fig. 2 - Weight-length relationship from fresh fish. $W_t (g) = 0.00819 \times L_f^{3.11} (cm)$; $n = 1934$; $r^2 = 0.988$.




Fig. 3 - Weight-length relationship from frozen fish. $W_t (g) = 0.00957 \times L_f^{3.06} (cm)$; $n = 419$; $r^2 = 0.988$.

Table 1. t test for significance of regression coefficients (weight-length regressions using fresh and frozen samples).

SAMPLES	n	Res. DF	b	S (b ₁ - b ₂)	t	p
(1) Fresh	1934	1932	3.0629			
(2) Frozen	419	417	3.1108	0.0176	2.7216	<0.01*

* Null hypothesis rejected ($p < 0.05$)

$W_t (g) = 0.00819 \times L_f^{3.11} (cm)$; $n = 1934$; $r^2 = 0.988$ for fresh fish (Figure 2), and:

$W_t (g) = 0.00957 \times L_f^{3.06} (cm)$; $n = 419$; $r^2 = 0.988$ for frozen fish (Figure 3).

Comparison between the slopes of the logarithmic transformation of the relations (Table 1) showed that they are significantly different as $p < 0.01$.

For $\alpha = 0.05$ we had to reject $H_0 : \beta_1 = \beta_2$, then $H_A : \beta_1 \neq \beta_2$.

Thus, for the present paper, only data obtained with fresh fish will be considered.

Total length-fork length relationship

The following relationship compares total

length (L_t) with fork length (L_f) (Figure 4):

$L_t (cm) = -0.447 + 1.137 \times L_f (cm)$; $n = 1880$; $r^2 = 0.994$.

Comparison between left and right otoliths

The t test was also applied to compare the regressions obtained from left and right otolith measurements (Table 2).

Null hypothesis was tested for all relations, for a significance level of 5%, and was not rejected, meaning that either could be used.

Regressions obtained between left otoliths (l_o) measurements (L =length, H =height and W =weight) and fork length (L_f) are:

$L_{l_o} (mm) = 1.759 + 0.224 \times L_f (cm)$; $n = 261$; $r^2 = 0.964$ (Figure 5)




Fig. 4 - Relation between total length and fork length. L_t (cm) = $-0.447 + 1.137 L_f$ (cm); $n = 1880$; $r^2 = 0.994$.




Fig. 5 - Relation between left otolith length and fork length. L_{lo} (mm) = $1.759 + 0.224 L_f$ (cm); $n = 261$; $r^2 = 0.964$.

$$H_{lo} \text{ (mm)} = 1.180 + 0.089 \times L_f \text{ (cm)}; n = 261; r^2 = 0.951 \text{ (Figure 6)}$$

$$W_{lo} \text{ (g)} = 0.0000278 \times L_f^{2.072} \text{ (cm)}; n = 261; r^2 = 0.962 \text{ (Figure 7)}$$

Age-length key by means of otoliths age reading

Age reading of whole otoliths proved to be quite impossible due to their opaqueness. Otolith

sections seemed to be more reliable structures for age determination in oceanic horse mackerel, although very thin sections did not present enough contrast between summer and winter zones, which are less differentiated in these temperate waters. Even with thick sections some opaque borders were very difficult to identify.

Table 2. *t* test for significance of regression coefficients (regressions between length, height and weight of left and right otoliths and the respective fork length).

OTOLITHS	n	Res. DF	b	S (b ₁ - b ₂)	t	p
Length						
(1) left otolith	261	259	0.2239	0.0043	0.2558	> 0.50 *
(2) right otolith	257	255	0.2228			
Height						
(1) left otolith	261	259	0.0887	0.0029	0.7855	> 0.40 *
(2) right otolith	257	255	0.0910			
Weight						
(1) left otolith	261	259	-2.0715	0.0407	0.3538	> 0.50 *
(2) right otolith	271	269	-2.0571			

* Null hypothesis not rejected ($p > 0.05$).




Fig. 6 - Relation between left otolith height and fork length. $H_{lo} \text{ (mm)} = 1.180 + 0.089 L_f \text{ (cm)}$; $n = 261$; $r^2 = 0.951$.




Fig. 7 - Relation between left otolith weight and fork length. $W_{lo} \text{ (g)} = 0.0000278 L_f^{2.072} \text{ (cm)}$; $n = 261$; $r^2 = 0.962$.

An age-length key and the mean fork length per age class are presented in Table 3.

Von Bertalanffy growth equation

Von Bertalanffy growth parameters were determined for both sexes together. Due to insufficient data ages 0-1 and 8-9 have been discarded for the calculations.

The values obtained using the "Ford-Walford plot" were $L_{\infty} = 49.8 \text{ cm}$ and $k = 0.23 \text{ year}^{-1}$; $t_0 = -0.23 \text{ year}$, was calculated as the mean of the t_0 values obtained by the substitution of the above mentioned values in the von Bertalanffy equation.

Using "Allen method" the parameters obtained were $L_{\infty} = 57.0 \text{ cm}$; $k = 0.17 \text{ year}^{-1}$ and $t_0 = -0.41 \text{ year}$.

The "Allen method" was also applied to try a better adjustment of the von Bertalanffy growth equation by giving t_0 a certain value. Thus, for $t_0 = -0.23$, $L_{\infty} = 52.9 \text{ cm}$ and $k = 0.20 \text{ year}^{-1}$, which seemed to be the values that gave a better fit of von Bertalanffy growth curve (Figure 8).

DISCUSSION

The relationships obtained between otolith measurements and fork length will eventually allow the determination of a prey size from otoliths found in stomach contents (e.g. *T.*




Figure 8. Von Bertalanffy growth equation for both sexes. $L_f \text{ (cm)} = 52.9 (1 - e^{-0.20(t + 0.23)})$; $n = 516$.

picturatus is the most important food organism for *Loligo forbesi* in the Azores (MARTINS 1982)).

Age determination using otolith sections was found to be the best method for obtaining growth parameters. Nevertheless some difficulties remained concerning the identification of year rings in fishes 3 or 4 years and up. Maximum age given in previous studies is 11 years (POLONSKI 1967) which seems to be compatible with our results indicating a maximum of 9 years of age.

Estimates of $L_{\infty} = 51.06 \text{ cm}$; $k = 0.14 \text{ year}^{-1}$;

Table 3. Age-length key obtained from otoliths reading.

Fork length class (cm)	Age (year)									
	0	1	2	3	4	5	6	7	8	9
10										
11										
12	17	1								
13	11	4								
14	22	5								
15	19	20	4							
16	9	23	6							
17	11	22	13							
18	2	14	16							
19		5	16	2						
20		2	17	2						
21		2	9	4						
22			8	8						
23			1	5	1					
24			5	7	3					
25			4	10	3					
26			1	9	4					
27			1	6	4					
28			1	6	7					
29				3	3					
30				1	8		1			
31				2	8	2				
32				3	9	2	1			
33				1	5	3				
34					3	2	1			
35					1	7	1			
36					2	3	2			
37					1	5	9			
38						10	11	1		
39						5	13	3		
40						1	2		1	1
41						1	2	1		
42							1	1		
43								1		
44										
45										
46										
47										
48										
49										
50										
n	91	98	102	69	62	41	44	7	1	1
Mean L _f	14.86	16.88	20.09	25.76	30.37	36.77	38.34	40.79	40.50	40.50

$t_0 = -1.58$ year, for total length, were calculated by Westhaus-Ekau and Ekau (personal communication) from a sample with a higher number of juveniles. Nevertheless our estimates, especially for k , seem to be more appropriate to this pelagic species as it seems to have a fast growth. For a better fit of our growth curve (Figure 8) more individuals under 15 cm and above 40 cm (fork length) should be sampled and otolith age determination should be reviewed. Otherwise Gompertz growth equation (RICKER 1980) should be applied.

ACKNOWLEDGEMENTS

Thanks are due to my husband and colleague Eduardo Isidro for using his computer statistical program, and to João Pereira for helping in data computation. Thanks are also due to Dr Ron O'Dor for reviewing the English text.

REFERENCES

- ALBUQUERQUE, R. M. 1954-56. Peixes de Portugal e Ilhas Adjacentes. Chaves para a sua determinação. - *Portugaliae Acta Biologica*. Série B, 5: 1-1164.
- ALLEN, K. R. 1985. Selected computer programs in FORTRAN for fish stock assessment. - *FAO Fisheries Technical Papers* 259: 129-138.
- BEDFORD, B. C. 1977. Further development of the technique of preparing thin sections of otoliths set in black polyester resin. - *International Council for the Exploration of the Sea* CM 1977 / F: 24. 11pp. (mimeo).
- BEDFORD, B. C. 1983. A method for preparing sections of large numbers of otoliths embedded in black polyester resin. - *Journal du Conseil Internationale pour l'Exploration de la Mer* 41: 4-12.
- CHRISTENSEN, J. M. 1964. Burning of otoliths, a technique for age determination of soles and other fish. - *Journal du Conseil Permanent Internationale pour l'Exploration de la Mer* 29: 73-81.
- FOWLER, H. W. 1936. The marine fishes of West Africa. - *Bulletin of the American Museum of Natural History* 70 (2): 1-688.
- HUREAU, J. C. & Th. MONOD (Eds). 1979. *Check-list of the Fishes of the North-Eastern Atlantic and of the Mediterranean*. Vols. I and II. Unesco, France.
- LETACONNOUX, R. 1951. Contribution à l'étude des espèces du genre *Trachurus* et spécialement du *Trachurus trachurus*. - *Mémoires. Office Scientifique et Technique des Pêches Maritimes* 15: 1-67.
- MARTINS, H. R. 1982. Biological studies of the exploited stock of *Loligo forbesi* (Mollusca: Cephalopoda) in the Azores. - *Journal of the Marine Biological Association of the United Kingdom* 62: 799-808.
- POLONSKII, A. S. 1967. Distribution of age and size composition of horse mackerel in the English Channel, the Celtic and North Seas 1963 to 1965. - *Annales Biologiques* 22: 182-183.
- RICKER, W. E. 1980. Calcul et interprétation des statistiques biologiques des populations de poissons. - *Bulletin of the Fisheries Research Board of Canada* 191F: 1-409.
- RYTOV, A. N. & T. M. RAZUMOVSKAYA 1984. A new subspecies of ocean scad, *Trachurus picturatus aleevi*, from the Southwestern Indian Ocean. - *Journal of Ichthyology* 24 (2): 1-6.
- SHABONEYEV, I. YE. 1973. Materials on the biology of horse mackerel *Trachurus picturatus picturatus* (Bowdich) in the eastern part of central Atlantic. - *Trudy Vsesoyuznogo Nauchnogo Issledovatel'skogo Instituta Morskogo Rybnogo Khozyaystva i Okeanografii. Moskva. (VNIRO)* 93: 91-100. (Translated from Russian by Jürgen Ropking, 1979.)
- SHABONEYEV, I. YE. & YE. I. RYAZANTSEVA 1977. Population structure of the oceanic horse mackerel (*Trachurus picturatus picturatus*). - *Journal of Ichthyology* 17 (6): 954-958.
- SHABONEYEV, I. YE. & A. N. KOTLYAR 1979. A comparative morphoecological analysis of the Eastern Pacific forms of *Trachurus symmetricus* and the Atlantic oceanic horse mackerel *Trachurus picturatus picturatus*. - *Journal of Ichthyology*, 19 (2): 24-29.
- SMITH-VANIZ, W. F. 1986. Carangidae. - Pp.

- 815-844 in: WHITEHEAD, P. J. P., M.-L. BAUCHOT, J.-C. HUREAU, J. NIELSEN and E. TORTONESE (Eds). *Fishes of the North-eastern Atlantic and the Mediterranean*. Volume II. Unesco, Paris.
- WILLIAMS, T. & B. C. BEDFORD 1974. The use of otoliths for age determination. - Pp. 114-123 in: BAGENAL, T. B. (Ed.) *Ageing of fish*. Proceedings of an International Symposium. Unwin Brothers Limited, England. 234pp..
- ZAR, J. H. 1974. *Biostatistical Analysis*. Prentice-Hall, Inc., N.J., 620pp.

Accepted 27 November 1989.