

Antioxidant activity evaluation from *Artemisia gorgonum* extracts

Neidy Rodrigues,^a Diana C. G. A. Pinto,^b Ana M. L. Seca,^{b,c} Helena Silva^d and Maria de Lourdes Pereira^a

^aDepartment of Biology & CICECO, University of Aveiro, 3810-193 Aveiro, Portugal

^bDepartment of Chemistry & QOPNA, University of Aveiro, 3810-193 Aveiro, Portugal

^cDCTD, University of Azores, 9501-801 Ponta Delgada, Portugal

^dDepartamento de Biology & CESAM, University of Aveiro, 3810-193 Aveiro, Portugal




Introduction

Artemisia gorgonum (Asteraceae) known as “Iosna or Iorna” (fig. 1) is used in Cape Verde in traditional medicine to treat inflammation, fever and gastroenteritis.^[1] The sesquiterpene lactone ridentin 1, furofuran lignan sesamin 2 and the flavonoid artemetin 3 (fig.2), isolated from *A. gorgonum* showed anti-plasmodium *in vitro* activity.^[2,3]



Recently, sesquiterpene lactones (seco-guaianolides) isolated from this plant, showed higher phytotoxic activity, and the authors suggested that they can be used as inspiration to develop new herbicides.^[4]

A few years ago was established that *A. gorgonum* volatile oil displays several biological properties including outstanding antioxidant activity.^[5]



However, to our best knowledge, no study on the potential antioxidant of other *A. gorgonum* extracts has been published.

Material & Methods

Plant collection and extracts preparation

Leaves of *A. gorgonum* were collected in Serra Malagueta Natural Park, Cape Verde, Santiago Island, in January 2012.

Four portion (500 mg each) of dried and powdered leaves of *A. gorgonum* were extracted with 20 mL of chloroform, methanol-chloroform (1:1), methanol and ethanol-water (7:3), C for during 30 minutes at 70° and then maintained at room temperature for 24 hours.^[6]

Antioxidant Activity


❖ Antioxidant activity was assayed by the DPPH (2,2-diphenyl-1-picrylhydrazyl (DPPH)) radical scavenging method.^[7] Briefly, to different concentration of ethanolic solutions of each extracts were added fixed volume of DPPH ethanolic solution and solvent (ethanol) to obtain in each case a fixed total volume. In each assay, a control was prepared, in which the sample or standard (quercetin and BHT) was substituted by the same amount of solvent.

❖ The absorbance of each solution was measured at 517 nm against a corresponding blank (ethanol solution) after 30 min. in dark at room temperature. The percentage of DPPH inhibition was calculated as follows

$$\% \text{ DPPH Inhibition} = \frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \times 100$$

Results and Discussion

- ❖ The extracts antioxidant activity was evaluated using the DPPH assay. Quercetin and BHT (butylated hydroxytoluene) were used as positive control.
- ❖ To our knowledge, this is the first study providing data on antioxidant activities of the Cape Verde *A. gorgonum* extracts.



- ❖ The plant extracts exhibit different EC₅₀ and in all cases higher than the standard compounds;
- ❖ In the case of the chloroform extract was not possible to obtain the EC₅₀ and the methanol-chloroform extract showed weak potential antioxidant as can be inferred from the EC₅₀ obtained;
- ❖ The methanolic extract presented the higher radical scavenging activity, although much higher than the standard compounds (BHT and quercetin);
- ❖ The results obtained suggest that *A. gorgonum* can be a potential source of natural antioxidant compounds;
- ❖ Chemical composition of the most active extracts will be studied and hopefully the obtained natural compounds will enlighten the extract antioxidant properties.

References

1. Gomes, I.; Gomes, S.; Vera-Cruz, M. T.; Leyens, T.; Kilian, N.; Lobin, W. Plantas endêmicas e árvores indígenas de Cabo Verde. DGA/ (GTZ) GmbH e “ Conservação da Biodiversidade”- DGA/UNDP-GEF CVI/00/G41 **2003**, p.12.
2. Ortet, R.; Prado, S.; Mouray, E.; Thomas, O. P. *Phytochemistry* **2008**, *69*, 2965.
3. Ortet, R.; Prado, S.; Regalado, E. L.; Valeriote, F. A.; Media, J.; Mendiola, J.; Thomas, O. P. *J. Ethnopharmacol.* **2011**, *138*, 637.
4. Macías, F. A.; Santana, A.; Yamahata, A.; Varela, R. M.; Fronczek, F. R.; Molinillo, J. M. G. *J. Nat. Prod.* **2012**, *75*, 1973.
5. Ortet, R.; Thomas, O. P.; Regalado, E. L.; Pino, J. A.; Filippi, J.-J.; Fernández, M. D. *Chem. Biodivers.* **2010**, *7*, 1332.
6. Smith, J. E., Tucker, D., Watson, K., Jones, G. L.,... *J. Ethnopharmacol* **2007**, *112*, 93.
7. Barreto, M. C.; Mendonça, E.; Gouveia, V.; Anjos, C.; Medeiros, J. S.; Seca, A. M. L.; Neto, A. I. *Arquipelago Life Mar. Sci.* **2012**, *29*, 58

Acknowledgments

Thanks are due to the University of Aveiro, Portuguese Foundation for Science and Technology (FCT), European Union, QREN, FEDER and COMPETE for funding the QOPNA research unit (project PEst-C/QUI/UI0062/2011), the CICECO (PEst-C/CTM/LA0011/2011-FCT) and CESAM.