



Dalila Carla Barbosa Sebastião

**Effect of temperature and prey in the biology of *Scymnus subvillosus* (Goeze)**  
**(Coleoptera: Coccinellidae)**

Departamento de Biologia

Ponta Delgada

2013



Dalila Carla Barbosa Sebastião

**Effect of temperature and prey in the biology of *Scymnus subvillosus* (Goeze)**  
**(Coleoptera: Coccinellidae)**

Dissertação para obtenção de Grau de Mestre em Biotecnologia em Controlo Biológico

Orientadores: Doutor António Onofre Costa Miranda Soares

Doutora Isabel Marisa Mateus Borges

Departamento de Biologia

Ponta Delgada

2013

## INDEX

ACKNOWLEDGEMENTS.....	4
ABSTRACT.....	5
RESUMO.....	7
1. Introduction .....	9
2. Material and Methods .....	11
2.1 The insects.....	11
2.2 Temperature and prey-dependence for development of the immature stages of <i>S. subvillosus</i> .....	11
2.3 Prey suitability of <i>A. fabae</i> , <i>M. donacis</i> and <i>M. persicae</i> for <i>S. subvillosus</i> .....	13
2.4 Voracity and nutritional physiology of the 4 <sup>th</sup> larval stage of <i>S. subvillosus</i> fed on <i>A. fabae</i> .....	14
2.5 Statistical analysis .....	15
3. Results .....	16
3.1 Temperature and prey-dependence for development of the immature stages of <i>S. subvillosus</i> .....	16
3.2 Prey suitability of <i>M. persicae</i> , <i>A. fabae</i> and <i>M. donacis</i> for <i>S. subvillosus</i> .....	20
3.3 Voracity and nutritional physiology of the 4 <sup>th</sup> larval stage of <i>S. subvillosus</i> fed on <i>A. fabae</i> .....	23
4. Discussion .....	24
5. General Conclusions .....	27
6. References .....	28

## ACKNOWLEDGEMENTS

O meu profundo e sentido agradecimento a todas as pessoas que contribuíram para a concretização desta dissertação, estimulando-me intelectualmente e emocionalmente.

- ✓ Ao Doutor António Onofre Soares, meu orientador, expresso o meu profundo agradecimento pela orientação e apoio incondicional que muito elevaram os meus conhecimentos científicos, pelo apoio, pelas opiniões e críticas, pela dedicação, pela total disponibilidade, pela sua amizade, boa disposição, simpatia, confiança que em mim depositou, pela partilha do saber e as valiosas contribuições para a realização do trabalho;
- ✓ À Doutora Isabel Borges, minha orientadora, expresso o meu profundo agradecimento pela sua total disponibilidade (mesmo de licença de maternidade), pela sua simpatia, apoio, pelo saber que transmitiu, atenção dispensada, pelo rigor, pela indispensável ajuda no trabalho de campo, pela colaboração do tratamento estatístico, dedicação, boa disposição e pela sua amizade e por todas contribuições que permitiram concretização do trabalho;
- ✓ À Natália Cabral, pela sua grande amizade, simpatia, palavras de incentivo, disponibilidade, pela preciosa e indispensável ajuda no trabalho de campo e assistência no laboratório;
- ✓ À Marta Vale, Lurdes Borges e Carolina Parelho, pela grande amizade, simpatia, disponibilidade por me apoiarem sempre com palavras positivas para animar o espírito;
- ✓ À Sofia e Sara Pereira pela excelente amizade, pelas palavras de incentivo, apoio e pelos momentos de descontração;
- ✓ Ao meu namorado, pelo carinho, paciência nos meus dias difíceis, compreensão e encorajamento, pela total disponibilidade de me acompanhar todos fins-de-semana para trabalho de laboratório durante todo este período, pela sua amizade e por tudo o que representa para mim;
- ✓ Aos meus Pais dirijo um agradecimento muito especial, por serem modelos de coragem, pelo apoio incondicional, pelas palavras de incentivo, dedicação, compreensão nos meus dias difíceis, pelo amor e carinho demonstrados e total ajuda na superação dos obstáculos que surgiram ao longo desta caminhada. A eles dedico este trabalho! LOVE YOU!

## ABSTRACT

*Scymnus subvillosus* (Goeze) is an aphidophagous predator present in the Azores Archipelago (Portugal), but it occurs at low densities. Recently it has been observed exploiting *Melanaphis donacis* (Passerini) on *Arundo donax* L. The *Scymnus* spp., although less studied, have, in recent years, been the subject of research in the perspective of their use as biological control agents.

In this study we aimed to determine the concomitant effects of four constant temperatures (15°C, 20°C, 25°C and 30°C) and three aphid species on survival and development of immature stages of *S. subvillosus*, to access the ecophysiological suitability of *M. persicae*, *A. fabae* and *M. donacis* and finely the voracity and nutritional physiology of the 4<sup>th</sup> larval stage fed on *A. fabae*.

The development time from larvae to adult decreased with increasing temperature, ranging from 61.5 days at 15°C to 10.4 days at 30°C. To complete pre-imaginal development on *M. donacis*, LDT was estimated to be 11.7°C and SET 196.3 °D. At 15°C, larvae failed to develop when fed on *A. fabae* and *M. persicae* but on *M. donacis* 22% of larvae have survived. . We also found that development time of immature stages were also prey-dependent, being *M. persicae* the least suitable.

Reproductive parameters were prey dependent, being roughly better with *A. fabae*.

Twelve hours 4<sup>th</sup> larval stage of *S. subvillosus* ingested 2.08 mg per day of biomass corresponding to an average of 10.5 aphids, allowing for a daily mean weight gain of 0.71 mg. The conversion efficiency and relative growth rate obtained was approximately 21% and 48% respectively.

The results obtained in the present study suggest that both, *A. fabae* and *M. donacis* are more suitable preys for development and reproduction of *S. subvillosus* than *M. persicae*.

Key-words: *S. subvillosus*, *A. fabae*, *M. donacis*, *M. persicae*, development parameters, reproductive parameters, nutritional physiology.

## RESUMO

*Scymnus subvillosus* (Goeze) é um predador afidífono presente no Arquipélago dos Açores (Portugal), que ocorre em baixas densidades., recentemente, foi observado a explorar *Melanaphis donacis* (Passerini) em *Arundo donax* L. O *Scymnus* spp., embora menos estudadas, têm sido nos últimos anos, alvo de investigação na perspectiva da sua utilização como agentes de controlo biológico.

Este estudo teve como objetivo determinar os efeitos concomitantes de quatro temperaturas constantes (15° C, 20 °C, 25 C e 30°C) e de três espécies de afídeos na sobrevivência e desenvolvimento dos estados imaturos de *S. subvillosus* , para obter a adequação ecofisiológica de *M. persicae*, *A. fabae* e *M. donacis* e finalmente a voracidade e fisiologia nutricional da fase larval L4 alimentadas com *A. fabae* .

O tempo de desenvolvimento a partir de larvas a adulto diminuiu com o aumento da temperatura, variando de 61,5 dias a 15 ° C para 10,4 dias a 30°C. Para completar o desenvolvimento pré -imaginal com *M. donacis*, LDT foi calculado obtendo o valor de 11,7°C e SET 196,3°D. A 15°C, as larvas não se desenvolveram quando alimentadas com *A. fabae* e *M. persicae*, mas com *M. donacis* 22 % das larvas sobreviveram. Assim sendo, o tempo de desenvolvimento dos imaturos também foi dependente da presa, sendo *M. persicae* a menos adequada.

Os parâmetros reprodutivos foram dependentes da presa, sendo melhor com *A. fabae*.

Larvas de *S. subvillosus* com doze horas no estado L<sub>4</sub>, ingeriram 2,08 mg por dia de biomassa, correspondente a uma média de 10,5 afídeos, permitindo um ganho de peso médio diário de 0,71 mg. A eficiência de conversão e a taxa de crescimento relativa foi de aproximadamente 21 % e 48 %, respectivamente.

Os resultados obtidos no presente estudo sugerem que ambas, *A. fabae* e *M. donacis* são presas mais adequadas para o desenvolvimento e reprodução de *S. subvillosus* do que *M. persicae*.

**Palavras-chave:** *S. subvillosus*, *A. fabae*, *M. donacis*, *M. persicae*, parâmetros de desenvolvimento, parâmetros reprodutivos e fisiologia nutricional.



# 1. INTRODUCCION

Ladybird predators (Coleoptera: Coccinellidae) are known as important biological control agents, especially against aphid populations (Hodek, van Emden and Honěk, 2012). Several species are used in biological control programs under its various strategies: classical, augmentative, inundative, in different crops and cultivation systems, either outdoors and greenhouses (Hodek and Michaud, 2008; Cabral et al., 2009). These predators have been extensively studied due to their biological, ecological and behavioral characteristics such as polyphagy, high voracity and rapid numerical and aggregative response (Hodek and Honěk, 1996; Obrycki and Kring, 1998; Dixon, 2000). Indeed the first case of successful biological control program was achieved with a ladybird species, *Rodolia cardinalis* (Mulsant) against the cottony cushion scale *Icerya purchasi* Maskell on citrus plantation in California (USA) (Dixon, 2000).

Among ladybirds the small *Scymnus* sp., although less studied, have, in recent years, been the subject of research in the perspective of their use as biological control agents (Uygun and Atlihan, 2000; Wanntorp, 2004; Pluke *et al.*, 2005; Woin et al., 2006). Comparing with larger species, the *Scymnus* spp., due to its small size and lower voracity, will be apparently less competitive and efficient as biological control agents. However, persisting at low densities of prey and presenting high longevities (Tawfik et al., 1973, Borges et al., 2011, 2013), these small ladybirds may be able to explore aphid colonies on earlier and /or later stages (Agarwala and Yasuda, 2001) and for a longer period of time, which would give them the possibility to exploit different spatial and temporal niches other than the larger ladybirds.

*Scymnus subvillosus* (Goeze) is an aphidophagous predator recorded for the Azores Archipelago (Portugal) (Soares et al. 2003; 2006), but occurring at low densities. It has been observed exploiting *Melanaphis donacis* (Passerini) on *Arundo*

*donax* L. This coccinellid is a cosmopolitan species, found in France and Mediterranean regions, Madeira archipelago (Portugal), Caucasus, Siberia, Minor Asia, North Africa and Germany (Gourreau, 1974; Raimundo and Alves, 1986). In the Portuguese mainland, it is common in arboreal agro-ecosystems, including apple orchards, orange, peach, plum, walnut and oak, where it exploits *Hyalopterus pruni* (Geoffroy) (Homoptera: Aphididae) (Raimundo and Alves, 1986). In Turkey, *S. subvillosus* is widespread (Atlihan and Güldal, 2009), and it is an important natural enemy against 15 aphid species (Aslan and Uygun, 2005). There it is common in agro-ecosystems and natural habitats as well, including stone fruits especially in *Prunus* spp. orchards, *Mallus* spp., and *Populus* spp., on apple, plum, cherry, pine and alder (Aslan and Uygun 2005).

The temperature and the nutritional quality of prey are decisive factors in the biological performance of insect predators, altering for instance the development time of pre-imaginal stages, longevity and reproductive parameters of adults (e.g., fecundity, fertility), and by this way contributing to differential population growth (McCaffrey and Horsburgh 1986; Nijima et al., 1986; Gibson et al., 1992; Hodek, 1993; Mohaghegh et al., 2001; Skirvin and Fenlon, 2003; Kontodimas and Stathas, 2005; Michaud, 2005; Cabral et al., 2006; Jalali et al., 2010). The estimation of the biological characteristics of natural enemies, such as time of development and the thermal constant may contribute to select the most appropriate biocontrol agent to be used under certain environmental conditions (Perdikis and Lykouressis, 2002) as well as the best conditions either thermal or nutritional for mass production.

In this study we aimed i) to determine the concomitant effects of four constant temperatures (15°C, 20°C, 25°C and 30°C) and three aphid species on survival and development of immature stages of *S. subvillosus*, ii) to access the ecophysiological

suitability of *M. persicae*, *A. fabae* and *M. donacis* as prey to *S. subvillosus* and iii) the voracity and nutritional physiology of the 4<sup>th</sup> larval stage fed on *A. fabae*.

## 2. MATERIAL AND METHODS

### 2.1 The insects

*Scymnus subvillosus* adults were collected in the field, on *Arundo donax* L., in S. Miguel Island, Azores (Portugal). The predator was found foraging and reproducing on *Melanaphis donacis* (Passerini). Ladybird adults were collected and reared in the laboratory at  $25 \pm 1^\circ\text{C}$ ,  $75 \pm 5\%$  RH and a light regime 16L:8D under fluorescent lamps. The predator was fed on a mixed diet of *Myzus persicae* (Sulzer) and *Aphis fabae* (Scopoli) (Homoptera: Aphididae), provided *ad libitum*, and complemented with pollen and honey. The mixed diet was provided to avoid food adaptation (Rana et al., 2002) and also to supply a wider group of nutrients. The prey species were reared on *Vicia faba* L. major at  $15 \pm 1^\circ\text{C}$ ,  $75 \pm 5\%$  RH and a light regime 16L: 8D under fluorescent lamps.

### 2.2 Temperature and prey-dependence for development of the immature stages of *S. subvillosus*

To determine the effect of temperature and prey on the development of the immature stages of *S. subvillosus*, the development time from egg to adult emergence was followed at 12 experimental treatments, corresponding each one to the following temperature,  $15^\circ\text{C}$ ,  $20^\circ\text{C}$ ,  $25^\circ\text{C}$  and  $30^\circ\text{C}$  and prey regimes of *A. fabae*, *M. donacis* or

*M. persicae* (N = 30). Each larvae was kept individually in a plastic box (diameter: 5cm, height: 3 cm) and was fed once a day *ad libitum*, with single diets of *A. fabae*, *M. donacis* or *M. persicae*. All assays were performed at  $75 \pm 5\%$  RH, with a photoperiod of 16L: 8D. Immature stages were observed twice a day (10:00 am and 17:00 pm) to record moulting and adult emergence.

The larval development time and survival were determined for each diet at each tested temperature. With the data on development at the different temperatures (15°C, 20°C, 25°C and 30°C), the lower developmental threshold (LDT) was determined. The (LDT) is the temperature below which a particular stage of an animal cannot develop. The relationship between the inverse of the development time and the temperature is nearly linear (Honěk and Kocourek, 1990). According to this, the following linear relationship was considered (Mota et al., 2008):

$$1/DT = aT + b$$

Where DT corresponds to the development time of the pre-imaginal, which is inversely proportional to the temperature (T), and *a* and *b* are the regression parameters. The LDT corresponds to T, when no development occurs, that is, when  $1/DT = 0$  (Honěk and Kocourek, 1988).

Sum of effective temperatures (SET), is the thermal time T which is the number of day degrees (°D) above lower developmental threshold for completion of a developmental stage (Hodek, van Emden and Honěk, 2012).

The sum of effective temperatures (SET) was calculated according to the model (Mota et al., 2008):

$$SET = \frac{[DT_{15}(15 - LDT_{15})] + [DT_{20}(20 - LDT_{20})] + [DT_{25}(25 - LDT_{25})] + [DT_{30}(30 - LDT_{30})]}{4}$$

DT<sub>x</sub>- development time to temperature x

LDT<sub>x</sub>- lower developmental threshold to temperature x

### 2.3 Prey suitability of *A. fabae*, *M. donacis* and *M. persicae* for *S. subvillosus*

The suitability of *A. fabae*, *M. donacis* and *M. persicae* as prey for *S. subvillosus* was evaluated by determining development data such as development time of immature stages (N=30), survival rate, weight of females soon after emergence and reproductive parameters, e.g., pre-oviposition time, fecundity and fertility. To evaluate reproductive performance of adults, 10 couples were formed and followed for 20 days on each prey diet. Each couple was kept inside a plastic box (diameter: 5cm height: 3cm) and fed *ad libitum* with single diets of *A. fabae*, *M. donacis* or *M. persicae*. All females were weighed before to the tests for reproductive parameters. Couples were observed once a day to obtain data on pre-oviposition time, fecundity and fertility.

All assays were performed at 25± 1°C, 75±5% RH and a light regime 16L: 8D under fluorescent lamps.

## 2.4 Voracity and nutritional physiology of the 4<sup>th</sup> larval stage of *S. subvillosus* fed on *A. fabae*

The larvae used on this experiment were previously fed *ad libitum* with *A. fabae* and reared under the same abiotic conditions of the previous experiments until they reached the 4<sup>th</sup> larval stage. The experiments were performed on *A. fabae* because it was the prey on which the predator presented faster development, high fecundity and fertility. To determine aphid natural mortality and weight loss due to dehydration, a control test was performed. Ten aphids were placed inside a plastic box (N=15). At the beginning of the experiments as well as at the end of the experiment (8:00 am/8:00 pm) the weight of aphids was recorded to measure the weight loss due to dehydration ( $PW_d$ ) and the survival rate ( $S_r$ ) of the aphids.

The test of consumption was performed with 12 hours old 4<sup>th</sup> instar larvae. Then larvae were subjected to starvation for a period of 12 hours, followed by a 24 hours test where they were allowed to consume prey.

Because the aphid prey dehydrated considerably on a 24 hours period, the predator was provided with 10 aphids in two feeding periods (8:00 am and 8:00 pm): in a total of 20 aphids per day/larvae. In order to determine the biomass consumption, the aphids were weighed before and after the 24hour feeding period. The number of prey consumed and naturally dead was recorded. In the beginning and at the end of the test the prey biomass (PW) and larvae weight (LW) were also recorded.

Voracity was estimated as the number of preys entirely or partially eaten (P). The voracity (V), biomass ingested (BI), relative growth rate (RGR %) and conversion efficiency (CE %) was also calculated (Borges, 2008) :

$$V_{S.subvillosus} = P \times S_r$$

$$BI_{S.subvillosus} = \frac{PW_f - PW_i - PW_d}{PW_i}$$

$$RGR \% = \frac{LW_f - LW_i}{LW_i} \times 100$$

$$CE \% = \frac{LW_f - LW_i}{BC} \times 100$$

Where  $PW_i$  and  $PW_f$  are prey initial and final weights respectively,  $LW_i$  and  $LW_f$  are the larval initial and final weights respectively.

## 2.5 Statistical analysis

Data normality and variance homogeneity were evaluated by the Kolmogorov-Smirnov and Levene's tests respectively. When normality and homogeneity of variance were confirmed, the parametric test of ANOVA was used and for multiple comparisons the Tukey test was performed. When data was not normal, non-parametric Kruskal-Wallis test was used, with multiple comparisons made by the Mann-Whitney test with the correction of Bonferroni.

The Bonferroni correction is a multiple-comparison correction used when several dependent or independent statistical tests are being performed simultaneously. To perform a Bonferroni correction, the critical  $P$  value ( $\alpha$ ) is divided by the number of comparisons being made. In the present study, 3 diets tested, therefore the critical  $P$  value is 0.017 ( $\alpha/3$ ).

SPSS v. 15.0 was used to perform statistical analysis.

### 3. RESULTS

3.1 Temperature and prey-dependence for development of the immature stages of *S. subvillosus*

Our results showed that the immature stages development time of *S. subvillosus* decreased with increasing temperature, as expected within the range of temperatures tested (Tables I, II and III).

At 15°C, only a few couples laid eggs and most of them were infertile. Due to this fact, statistical analyses on developmental data did not include embryonic development. To overcome the reduced number of eggs obtained at 15°C, couples were maintained at 25°C to allow oviposition. The eggs were then collected and transferred to 15°C in order to obtain new born larvae to follow its development.

Lower development threshold LDT (°C) and sum effective temperatures SET (°D) for immature development of *S. subvillosus* fed on *M. donacis* range from 9.4°C for L1 to 12°C for L4 and SET from 16.7°D for L2 to 70.6°D for pupa (Table IV). To complete pre-imaginal development, LDT was estimated to be 11.7°C and SET 196.3°D (Table IV). The thermal parameters of larvae fed on *A. fabae*, *M. persicae* was not calculated because data from 3 temperatures (20°C, 25°C and 30°C) leads to an inaccurate and unrealistic estimates of the LDT and SET values.



**Table I:** Development time of immature stages (days  $\pm$  SE) of *S.subvillosus* under constant temperatures (15°C, 20°C, 25°C and 30°C). L<sub>1</sub>, L<sub>2</sub>, L<sub>3</sub>, L<sub>4</sub> and P: first, second, third and fourth larval stages and pupa, respectively fed with *M. persicae*.

Developmental stages	15°C	20°C	25°C	30°C	ANOVA (F) or Kruskal-Wallis ( $\chi^2$ )
L <sub>1</sub>	—	2.34 $\pm$ 0.15a*	2.03 $\pm$ 0.07ab	1.84 $\pm$ 0.08b	F = 3.84, df = 2, p = 0.029
L <sub>2</sub>	—	1.67 $\pm$ 0.15a	1.57 $\pm$ 0.21a	0.97 $\pm$ 0.60b	$\chi^2$ = 11.06, df = 2, p = 0.004
L <sub>3</sub>	—	1.85 $\pm$ 0.24a	1.69 $\pm$ 0.43ab	1.42 $\pm$ 0.53b	F = 3.36, df = 2, p = 0.046
L <sub>4</sub>	—	6.04 $\pm$ 0.14a	4.21 $\pm$ 0.65b	4.17 $\pm$ 0.44b	F = 47.17, df = 2, p $\leq$ 0.0001
P	—	7.77 $\pm$ 0.26a	4.14 $\pm$ 0.19b	3.86 $\pm$ 0.10b	F = 272.28, df = 2, p $\leq$ 0.0001
<b>Total</b>	—	19.5 $\pm$ 0.23a	13.8 $\pm$ 0.30b	12.3 $\pm$ 0.17c	F = 258.32, df = 2, p $\leq$ 0.0001

\* Means in each row followed by different letters significantly different at p < 0.05 (ANOVA or Kruskal-Wallis).

**Table II:** Development time of immature stages (days  $\pm$  SE) of *S.subvillosus* under constant temperatures (15°C, 20°C, 25°C and 30°C). L<sub>1</sub>, L<sub>2</sub>, L<sub>3</sub>, L<sub>4</sub> and P: first, second, third and fourth larval stages and pupa, respectively fed with *A. fabae*.

Developmental stages	15°C	20°C	25°C	30°C	ANOVA (F) or Kruskal-Wallis ( $\chi^2$ )
L <sub>1</sub>	—	2.15 $\pm$ 0.07a*	1.80 $\pm$ 0.06b	2.07 $\pm$ 0.04a	$\chi^2 = 11.07$ , df = 2, p $\leq$ 0.0001
L <sub>2</sub>	—	2.08 $\pm$ 0.04a	1.03 $\pm$ 0.09b	1.20 $\pm$ 0.08b	$\chi^2 = 39.4$ , df = 2, p $\leq$ 0.0001
L <sub>3</sub>	—	1.63 $\pm$ 0.10a	1.38 $\pm$ 0.12ab	1.08 $\pm$ 0.08b	F = 7.5, df = 2, p = 0.001
L <sub>4</sub>	—	5.11 $\pm$ 0.14a	4.22 $\pm$ 0.09b	3.75 $\pm$ 0.13c	F = 32.45, df = 2, p $\leq$ 0.0001
P	—	7.14 $\pm$ 0.08a	4.75 $\pm$ 0.07b	4.17 $\pm$ 0.08c	F = 411.89, df = 2, p $\leq$ 0.0001
<b>Total</b>	—	17.34 $\pm$ 0.14a	13.17 $\pm$ 0.12b	12.08 $\pm$ 0.20c	$\chi^2 = 11.07$ , df = 2, p $\leq$ 0.0001

\*Means in each row followed by different letters significantly different at p < 0.05 (ANOVA or Kruskal-Wallis).

**Table III:** Development time of immature stages (days  $\pm$  SE) of *S. subvillosus* under constant temperatures (15°C, 20°C, 25°C and 30°C). L<sub>1</sub>, L<sub>2</sub>, L<sub>3</sub>, L<sub>4</sub> and P: first, second, third and fourth larval stages and pupa, respectively fed with *M. donacis*.

Developmental stages	15°C	20°C	25°C	30°C	Kruskal-Wallis ( $\chi^2$ )
L <sub>1</sub>	7.11 $\pm$ 0.37a*	2.95 $\pm$ 0.06b	1.97 $\pm$ 0.05c	1.77 $\pm$ 0.07c	$\chi^2 = 76.66$ , df = 3, p $\leq$ 0.0001
L <sub>2</sub>	6.01 $\pm$ 0.28a	1.81 $\pm$ 0.14b	1.22 $\pm$ 0.08c	0.82 $\pm$ 0.08d	$\chi^2 = 35.54$ , df = 3, p $\leq$ 0.0001
L <sub>3</sub>	6.61 $\pm$ 0.37a	2.17 $\pm$ 0.12b	1.67 $\pm$ 0.12c	0.89 $\pm$ 0.07d	$\chi^2 = 31.61$ , df = 3, p $\leq$ 0.0001
L <sub>4</sub>	20.63 $\pm$ 0.45a	7.06 $\pm$ 0.15b	4.5 $\pm$ 0.13c	3.25 $\pm$ 0.19d	$\chi^2 = 28.96$ , df = 3, p $\leq$ 0.0001
P	23.31 $\pm$ 0.83a	8.0 $\pm$ 0.09b	5.63 $\pm$ 0.08c	3.75 $\pm$ 0.09d	$\chi^2 = 29.60$ , df = 3, p $\leq$ 0.0001
<b>Total</b>	61.50 $\pm$ 1.96a	22.00 $\pm$ 0.22b	14.75 $\pm$ 0.11c	10.50 $\pm$ 0.13d	$\chi^2 = 21.86$ , df = 3, p $\leq$ 0.0001

\* Means in each row followed by different letters significantly different at p < 0.05 (Kruskal-Wallis).

**Table IV:** Lower development threshold (LDT) and sum of effective temperatures (SET  $\pm$  SE) required for the development of the immature stages of *S. subvillosus* fed on *M. donacis*, with the respective coefficient of determination ( $R^2$ ), slope (a) and intercept (b). L1, L2, L3, L4 and P: first, second, third and fourth larval stages and pupa, respectively.


Stages	a	b	$R^2$	LDT ( $^{\circ}$ C)	SET ( $^{\circ}$ D)
L1	0.0301	-0.2825	0.9687	9.4	34.2 $\pm$ 2.13
L2	0.0629	-0.7302	0.9766	11.6	16.7 $\pm$ 1.40
L3	0.0512	-0.5962	0.9901	11.6	20.1 $\pm$ 0.97
L4	0.0171	-0.2046	0.9983	12.0	59.1 $\pm$ 1.19
P	0.0143	-0.1693	0.993	11.8	70.6 $\pm$ 2.18
<b>Total</b>	0.0052	0.0606	0.9949	11.7	196.3 $\pm$ 5.93

### 3.2 Prey suitability of *M. persicae*, *A. fabae* and *M. donacis* for *S. subvillosus*

**Table V:** Development time of immature stages (Days  $\pm$  SE) of *S. subvillosus* fed with 3 different preys (different letters mean significant differences followed in each row).


Temperature	<i>M. persicae</i>	<i>A. fabae</i>	<i>M. donacis</i>	Kruskal-Wallis ( $\chi^2$ )
15 $^{\circ}$ C	-	-	61.5 $\pm$ 1.96	-
20 $^{\circ}$ C	20.3 $\pm$ 0.14a	17.5 $\pm$ 0.31b	21.6 $\pm$ 0.15c	$\chi^2 = 48.43$ , df = 2, p $\leq$ 0.0001
25 $^{\circ}$ C	13.8 $\pm$ 0.26a	13.2 $\pm$ 0.12a	14.9 $\pm$ 0.04b	$\chi^2 = 26.54$ , df = 2, p $\leq$ 0.0001
30 $^{\circ}$ C	12.3 $\pm$ 0.17a	12.3 $\pm$ 0.25a	10.4 $\pm$ 0.10b	$\chi^2 = 23.01$ , df = 2, p $\leq$ 0.0001

At 20 $^{\circ}$ C, larvae of *S. subvillosus* developed significantly faster fed on *A. fabae* and slower on *M. donacis*. At 25 $^{\circ}$ C it developed significantly slower on *M. donacis* and significantly faster on that prey at 30 $^{\circ}$ C. (Table V).




**Figure 1:** Survival rate (mean  $\pm$  SE) of immature stages of *S. subvillosus* fed with different preys.

Concerning survival rates, we found that at 15 °C larvae failed to develop into adult when fed on *A. fabae* and *M. persicae* but on *M. donacis* 22% of larvae have survived. At 20 and 25°C larval survival was high, ranging from 74.1% to 85.7% (Figure 1).




**Figure 2:** Pre-oviposition time (Days  $\pm$  SE) of *S. subvillosus* females at 25°C fed with 3 different preys (different letters mean significant differences).

Pre-oviposition time was significantly longer when females fed on *M. persicae* and no significant differences were found for *A. fabae* and *M. donacis* (ANOVA,  $F = 17.95$ ,  $df = 2$ ,  $p \leq 0.0001$ ) (Figure 2).



**Figure 3:** Fecundity (mean  $\pm$  SE) of *S. subvillosus* at 25°C fed with *M. persicae*, *A. fabae* and *M. donacis*.

Fecundity did not significantly differ with prey species (ANOVA,  $F = 1.59$ ,  $df = 2$ ,  $p = 0.224$ ). Females of *S. subvillosus* laid on average a minimum of 71 eggs and a maximum of 119 eggs when fed on *M. persicae* and *A. fabae*, respectively. Despite the absence of significant differences in the mean number of eggs laid, we found that the proportion of ovipositing females was not the same e.g. 90%, 70% and 50% for *M. donacis*, *A. fabae* and *M. persicae*, respectively.



**Figure 4:** Percentage of fertility (mean  $\pm$  SE) *S. subvillosus* at 25°C.

The fertility of *S. subvillosus* eggs was high with prey *A. fabae* (86.7%) and lower on *M. persicae* (62%) but no significant differences were observed. (Kruskal-Wallis  $\chi^2 = 5.03$ ,  $df = 2$ ,  $p = 0.081$ ) (Figure 4).

### 3.3 Voracity and nutritional physiology of the 4th larval stage of *S. subvillosus* fed on *A. fabae*

**Table VI:** Feeding parameters of the 4<sup>th</sup> larval stage of *S. subvillosus*.

Feeding parameters	Mean $\pm$ SE
Larval weight initial (mg $\pm$ SE)	1.69 $\pm$ 0.08
Larval weight final (mg $\pm$ SE)	2.40 $\pm$ 0.06
Mean weight gain (mg $\pm$ SE)	0.71 $\pm$ 0.11
Voracity (number of aphids eaten $\pm$ SE)	10.5 $\pm$ 0.78
Voracity (mg of biomass ingested $\pm$ SE)	2.08 $\pm$ 0.29
Relative growth rate (%)	47.7 $\pm$ 1.93
Conversion efficiency (%)	21.2 $\pm$ 9.42

The mean weight gain by a 12 hours 4<sup>th</sup> larval instar was 0.71 mg, corresponding roughly to 50% of their initial body weight (1.69 mg) (Table VI). The 2.08 mg of aphid biomass ingested, correspond to approximately 10.5 aphids eaten, is converted to larvae biomass on a rate of 21% (Table VI).

## 4. DISCUSSION

Environmental temperature is one of the most important factors determining physiological process on poikilothermic species with direct implications on development rate and survival (Honěk and Kocourek, 1990; Chapman, 1998). Several authors have shown that food quality also affects the development rate and survival (Blackman, 1967; Obrycki and Orr, 1990; Kalushkov, 1998; Kalushkov and Hodek, 2001; Işikber and Copland, 2002), as well reproductive performance, such as fecundity and fertility (Blackman, 1967; Nijima et al., 1986; Hodek, 1993; Michaud, 2005).

As we expected, developmental time of the immature stages decreased with increasing temperature. Our results also indicated that developmental rate and survival were concomitantly temperature and prey-dependent, which is consistent with previous studies (Atlihan et al., 1999; Satar and Uygun 2012). Survival rate was higher when larvae fed on *M. donacis*. Developmental rate decrease at 20°C and 30°C feeding on *A. fabae* and *M. donacis*, respectively. The most paradigmatic example was the incapacity of *S. subvillosus* to complete the development at 15°C fed with *M. persicae* and *A. fabae*. The importance of prey quality on developmental rate could be revealed by differences by the sum of effective temperatures (SET). In our experiment, *S. subvillosus* required 196.3°D, but the same species requires 329°D and 286°D to develop from egg to adult when fed *H. pruni* and *A. gossypii*, respectively (Atlihan and Chi 2008; Satar and Uygun, 2012).

Growth and development of insect species occur only across specific range of temperature (Nedvěd and Honěk, 2012) and thus the thermal parameters as lower developmental threshold and thermal constant are useful indicators to predict its potential distribution and abundance (Messenger, 1970). The LDT of *S. subvillosus* was estimated as 11.7°C for total immature stages development fed on *M. donacis*. Satar and Uygun (2012) determined 10.26°C from egg to adult of *S. subvillosus* fed on *A. gossypii*. These values are



quite similar to other *Scymnus* species (*S. hofmanii*: 10.1°C, *Scymnus frontalis*: 11.7°C fed on *Diuraphis noxia* (Mordvilko); *S. syriacus*: 11.3°C; *S. levaillanti*: 11.7°C fed *A. gossypii*; (Kawauchi, 1985; Naranjo et al., 1990; Emami et al. 1998; Uygun and Atlihan, 2000). Atlihan and Chi (2008) estimated a lower LDT for *S. subvillosus* fed on *H. pruni*; 7.1°C. Some of the differences can be partially explained by the differences on regional acclimatization. Satar and Uygun (2012) collected their *S. subvillosus* in the Mediterranean region where mild winters occur whereas Atlihan and Chi (2008) collected individuals in Turkey (Van region) where there is a harsh and long winter condition. The combination of a high LDT and a low SET guarantees a fast development at high temperatures, in contrast to cold adapted species whose LDT is low and SET high (Trudgill, 1995).

Egg production requires nutritional intake beyond a maintenance level, and thus high quality food is important for supporting reproductive capacity (Seagraves, 2009). Our results showed that *M. donacis* and *A. fabae* are equally suitable preys and on the other hand *M. persicae* leads to worse biological performance. In the Scymninae the pre-oviposition period varies with temperature and prey suitability even among closely related species (Naranjo et al., 1990; Uygun and Atlihan, 2000; Nedvěd and Honěk, 2012).

From an ecophysiological point of view, the three prey species tested on our study can be considered as essential foods. Indeed, under the light of Hodek, van Emden and Honěk (2012) criteria, the consumption of those prey species provided growth and development of larvae and reproduction by adults. Other essential prey for *S. subvillosus* includes *Aphis sambuci*, *Hyalopterus pruni*, and *Aphis gossypii* are reported as essential foods for *S. subvillosus* (Klausnitzer, 1992; Atlihan and Chi, 2008; Atlihan and Guldal, 2009; Satar and Uygun, 2012). However, essential foods, show varying degrees of favorability, enabling different developmental rates, fecundity, and survival (Hodek, 1993; Hodek and Honek, 1996; Kalushkov, 1998; Kalushkov and Hodek, 2004; Soares et al.,

2005; Cabral et al., 2006). Given that, we consider that *A. fabae* and *M. donacis* are more suitable food resources than *M. persicae*. Those results could reflect the co-occurrence between *S. subvillosus* and the *A. fabae* and *M. donacis* typically found in Azorean costal habitats, contrarily to *M. persicae*. Indeed according to Rana, Dixon and Jarosik 2002, the prey specialization increase fitness on ladybirds.

Body-weight increase following predation is a good indicator of energy intake and associated costs (Frazer, 1988). The mean daily weight gain by the 4<sup>th</sup> larval stage of *S. subvillosus* during 24h was 0.71 mg after a daily consumption of 10.5 aphids, which correspond to a 2.08 mg of biomass ingested. The ladybird handled around 6.3 mg of biomass, but has ingested only 2.08 mg, which means that after spending energy on basal metabolism, excretion and others activities, she assimilates only about 33% of weight of their preys. The larvae increased for doubled their size, 21% of the biomass ingested (average of 0.44 mg) is converted to larvae biomass so close 79% was spent on basal metabolism and metabolic activities of the larvae. The relative growth rate was approximately of 48%. Borges (2008), obtained similar value for conversion efficiency for *S. nubilus*, was 26%. Our results show that the high mortality of the prey may be because the ladybird satiates quickly and eats little of each aphid and spends much energy looking for a better or new prey.

## 5. GENERAL CONCLUSIONS

Our results clearly show the concomitant effect of temperature and prey on developmental time and survival rate of immature stage of *S. subvillosus*. Least evident were the differences concerning fecundity and fertility.

This study indicates that *M. persicae*, *A. fabae* and *M. donacis* are essential prey species for *S. subvillosus*. However diets differ in suitability, with *M. donacis* and *A. fabae* being better than *M. persicae* but equally suitable. The higher survival rate was observed with *M. donacis* at 20 °C and 25°C.

According to our results and published data, optimal temperature for population growth of *S. subvillosus* and for its mass production could be somewhere between 20 to 25°C.

## 6. REFERENCES

- Agarwala, B. K., Yasuda, H., 2001. Larval interactions in aphidophagous predators: effectiveness of wax cover as defence shield of *Scymnus* larvae against predation from syrphids. *Entomologia Experimentalis et Applicata* 100, 101–107.
- Aslan, M., Uygun, N., 2005. The Aphidophagus Coccinellid (Coleoptera: Coccinellidae) Species in Kahramanmaraş, Turkey. *Turkish Journal Zoology* 29, 1–8.
- Atlihan, R., Denizhan, E., Yaşar, B., 1999. Effects of different preys on development and fecundity of *Scymnus subvillosus* Goeze (Coleoptera: Coccinellidae). Proc. IV Turkish National Congress of Biological Control (Adana, Turkey), 397–406 (Turkish, with English summary).
- Atlihan, R., Chi, H., 2008. Temperature- dependent development and demography of *Scymnus subvillosus* (Coleoptera: Coccinellidae) reared on *Hyalopterus pruni* (Homoptera: Aphididae). *Journal of Economic Entomology* 101, 325–333.
- Atlihan, R., Güldal, H., 2009. Prey density-dependent feeding activity and life history of *Scymnus subvillosus*. *Phytoparasitica* 37, 35–41.
- Blackman, R.L., 1967. The effects of different prey on *Adalia bipunctata* L. and *Coccinella 7-punctata* L. *Annals Applied. Biology* 59, 207–219.
- Borges, I.M.M., 2008. Life history evolution in aphidophagous and coccidophagous Coccinellidae (Coleoptera). PhD Thesis. University of the Azores, Portugal.
- Borges, I., Soares, A.O., Magro, A., Hemptinne, J.L., 2011. Prey availability in time and space is a driving force in life history evolution of predatory insects. *Evolutionary Ecology* 25, 1307–1319.

- Borges, I., Hemptinne, J.L., Soares, A.O., 2013. Contrasting population growth parameters of the aphidophagous *Scymnus nubilus* and the coccidophagous *Nephus reunioni*. *BioControl* 58, 351–357.
- Cabral, S., Soares, A.O., Moura, R., Garcia, P., 2006. Suitability of *Aphis fabae*, *Myzus persicae* (Homoptera: Aphididae) and *Aleyrodes proletella* (Homoptera: Aleyrodidae) as prey for *Coccinella undecimpunctata* (Coleoptera: Coccinellidae). *Biological Control* 39, 434–440.
- Cabral, S., Soares, A.O., Garcia, P., 2009. Predation by *Coccinella undecimpunctata* L. (Coleoptera: Coccinellidae) on *Myzus persicae* Sulzer (Homoptera: Aphididae): effect of prey density. *Biological Control* 50, 25–29.
- Chapman, R.F., 1998. *The insects: structure and function*, 4<sup>th</sup> Ed. Cambridge University Press, UK, pp.770.
- Dixon, A. F. G., 2000. *Insect predator–prey dynamics: ladybirds and biological control*. Cambridge University Press, Cambridge, United Kingdom.
- Emami, M. S., Sahragard, A., Haji-Zadeh J., 1998. Effect of different temperatures on the development of *Scymnus syriacus* (Coleoptera: Coccinellidae). *Applied Entomology and Phytopathology* 66, 21–22.
- Frazer, B.D. 1988. *Predators Aphids: Their Biology, Natural Enemies and Control* (ed. by AK Minks & P Harrewijn), Elsevier Science, Amsterdam, The Netherlands Vol. B, pp. 217–230.
- Gibson, R., Elliot, N., Schaefer, P., 1992. Life history and development of *Scymnus frontalis* (Fabricius) (Coleoptera: Coccinellidae) on four species of aphid. *Journal of the Kansas Entomology Society* 65, 410–415.

- Gourreau, J. M., 1974. *Systematique de la tribu des Scymnini (Coccinellidae)*. Institut National de la Recherche Agronomique, Paris.
- Hodek, I., 1993. Habitat and food specificity in aphidophagous predators. *Biocontrol Science and Tecnology* 3, 91–100.
- Hodek, I., Honek, A., 1996. *Ecology of Coccinellidae*. Kluwer Academic, Dordrecht, the Netherlands.
- Hodek, I., Michaud, J.P., 2008. Why is *Coccinella septempunctata* so successful? (a point-of-view). *European Journal of Entomology* 105, 1–12.
- Hodek, I., Van Emden, H.F., Honek, A., 2012. *Ecology and Behaviour of the Ladybird Beetles (Coccinellidae)*. Blackwell Publishing Ltd.
- Honek, A., Kocourek, F., 1988. Thermal requirements for development of aphidophagous Coccinellidae (Coleoptera), Chrysopidae, Hemerobiidae (Neuroptera), and Syrphidae (Diptera): some general trends. *Oecologia* 76, 455–460.
- Honek, A., Kocourek, F., 1990. Temperature and development time in insects: a general relationship between thermal constants. *Zoologische Jahrbücher Systematik* 117, 401–439.
- Işıkber, A.A., Copland, M.J.W., 2002. Effects of various aphid foods on *Cycloneda sanguinea*. *Entomologia Experimentalis et Applicata* 102, 93–97.
- Jalali, M.A., Tirry, L., De Clercq, P., 2010. Effect of temperature on the functional response of *Adalia bipunctata* to *Myzus persicae*. *Biocontrol* 55, 261–269.
- Kalushkov, P., 1998. Ten aphid species (Sternorrhyncha: Aphidae) as prey for *Adalia bipunctata* (Coleoptera: Coccinellidae). *European Journal of Entomology* 95, 343–349.

- Kalushkov, P., Hodek, I., 2001. New essential aphid prey for *Anatis ocellata* and *Calvia quatuordecimguttata* (Coleoptera: Coccinellidae). *Biocontrol Science Technology* 11, 35–39.
- Kalushkov, P., Hodek, I., 2004. The effects of thirteen species of aphids on some life history parameters of the ladybird *Coccinella septempunctata*. *BioControl* 49, 21–32.
- Klausnitzer, B., 1992. Coccinelliden als Prädatoren der Holunderblattlaus (*Aphis sambuci* L.) im Wärme Frühjahr 1992. *Entologische Nachrichten und Berichte* 36, 185–190.
- Kawauchi, S.E., 1985. The threshold temperature and thermal constant for development from the egg to the adult form of *Coccinella septempunctata brucki*, *Propylea japonica* and *Scymnus (Pullus) hoffmanni* (Coleoptera, Coccinellidae). *Kurume University Journal* 32, 45–51.
- Kontodimas, D.C., Stathas, G.J., 2005. Phenology, fecundity and life table parameters of the predator *Hippodamia variegata* reared on *Dysaphis crataegi*. *BioControl* 50, 223–233.
- McCaffrey, J.P., Horsburgh, R.L., 1986. Functional response of *Orius insidiosus* (Hemiptera: Anthocoridae) to the European red mite, *Panonychus ulmi* (Acari: Tetranychidae), at different constant temperatures. *Environmental Entomology* 15, 532–535.
- Messenger, P.S., 1970. Bioclimatic inputs to biological control and pest management programs. *Concepts of Pest Management* (Ed. By R.L. Rabb. & F.E. Guthrie) North Carolina State University Press, Raleigh, 84–99.
- Michaud, J.P., 2005. On the assessment of prey suitability in aphidophagous Coccinellidae. *European Journal of Entomology* 102, 385–390.
- Mota, A., Soares, A.O., Garcia, P., 2008. Temperature dependence for development of the whitefly predator *Clitostethus arcuatus* (Rossi). *BioControl* 53, 603–613.

- Mohaghegh, J., De Clercq, P., Tirry, L., 2001. Functional response of the predator *Podisus maculiventris* (Say) and *Podisus nigrispinus* (Dallas) (Het.: Pentatomidae) to the beet armyworm, *Spodoptera exigua* (Hübner) (Lep.: Noctuidae): effect of temperature. *Journal Applied Entomology* 125, 131–134.
- Naranjo, S.E., Gibson, R.L., Walgenbach, D.D., 1990. Development, survival and reproduction of *Scymnus frontalis* (Coleoptera: Coccinellidae), an imported predator of Russian wheat aphid, at four fluctuating temperatures. *Annals of the Entomological Society of America* 83, 527–531.
- Nedvěd, O., Honěk, A., 2012. Life history and development, In: Hodek, I., Van Emden, H.F., Honek, A., *Ecology and Behavior of the Ladybirds Beetles (Coccinellidae)*, Blackwell Publishing Ltd, pp. 54–109.
- Nijjima, K., Matsuka, M., Okada, I., 1986. Artificial diets for an aphidophagous coccinellid, *Harmonia axyridis*, and its nutrition (Minireview). *Ecology of Aphidophaga*. Academia, Prague.
- Obrycki, J.J. & Orr, C.J., 1990. Suitability of three prey species for Nearctic populations of *Coccinella septempunctata*, *Hippodamia variegata*, and *Propylea quatuordecimpunctata* (Coleoptera: Coccinellidae). *Journal Economic Entomology* 4, 1292–1297.
- Obrycki, J.J., Kring, T.T., 1998. Predaceous coccinellidae in biological control. *Annual Review of Entomology* 43, 295–321.
- Perdikis, D.C., Lykouressis, D.P., 2002. Thermal requirements of the polyphagous predator *Macrolophus pygmaeus* (Hemiptera: Miridae). *Environmental Entomology* 31, 661–667.



- Pluke, R.W.H., Escribano, A., Michaud, J.P., Stansly, P.A., 2005. Potential impact of lady beetles on *Diaphorina citri* (Homoptera: Psyllidae) in Puerto Rico. *Florida Entomologist* 88, 123–128.
- Raimundo, A.A.C., Alves, M.L.G., 1986. Revisão dos coccinélídeos de Portugal. Universidade de Évora, Évora.
- Rana, J.S., Dixon, A.F.G., Jarosik, V., 2002. Costs and benefits of prey specialization in a generalista insect predator. *Journal of Animal Ecology* 71, 15–22.
- Satar, G., Uygun, N., 2012. The effects of various temperatures on development and fecundity of *Scymnus subvillosus* (Goeze) (Coleoptera: Coccinellidae) feeding on *Aphis gossypii* Glover (Hemiptera: Aphididae). *Türk. biyo. müt. derg.*, 3, 169–182.
- Seagraves, M.P., 2009. Lady beetle oviposition behaviour in response to the tropic environment. *Biological Control* 51, 313–322.
- Skirvin, D.J., Fenlon, J.S., 2003. The effect of temperature on the functional response of *Phytoseiulus persimilis* (Acari: Phytoseiidae). *Experimental and Applied Acarology* 31, 37–49.
- Soares, A.O., Coderre, D., Schanderl, H., 2001. Fitness of two phenotypes of *Harmonia axyridis* (Coleoptera: Coccinellidae). *European Journal of Entomology* 98, 287–293.
- Soares, A.O., Elias, R.B., Resendes, R., Figueiredo, H., 2003. Contribution to the knowledge of the Coccinellidae (Coleoptera) fauna from the Azores islands. *Life and Marine Sciences* 20, 47–53.
- Soares, A.O., Coderre, D., Schanderl, H., 2004. Dietary self-selection behaviour by the adults of the aphidophagous ladybeetle *Harmonia axyridis* (Coleoptera: Coccinellidae). *Journal of Animal Ecology* 73, 478–486.

- Soares, A.O., Coderre, D., Schanderl, H., 2005. Influence of prey quality on the fitness of two phenotypes of *Harmonia axyridis* adults. *Entomologia Experimentalis et Applicata* 114, 227–232.
- Soares A.O., Borges, I, Cabral, S. Figueiredo, H., Resendes, R., 2006. New Records of Coccinellidae (Coleoptera) to the Azores islands.
- Tawfik, M.F.S., Abdul-Nasr, S., Saad, B. M., 1973. The biology of *Scymnus interruptus* Goeze (Coleoptera: Coccinellidae). *Bulletin de la Societe Entomologique d’Egypte* 57, 9–26.
- Turgill, D.L., 1995. Why do tropical poikilothermic organisms tend to have a higher threshold temperature for development than temperate ones. *Functional Ecology* 9, 136–137.
- Uygun, N., Atlıhan, R., 2000. The effect of temperature on development and fecundity of *Scymnus levaillanti*. *BioControl* 45, 453–462.
- Wanntorp, H.E., 2004. “Musical chairs”: the Swedish species of *Scymnus* subg. *Neopullus* (Coleoptera, Coccinellidae) change places. *Entomologisk Tidskrift* 125, 103–109.
- Woin, N., Volkmar, C., Ghogomu, T., 2006. Numerical response of predatory ladybirds (Coccinellidae) to aphid outbreaks and their diversity in major rice ecosystems of Cameroon. *Archives of Phytopathology and Plant Protection* 39, 189–196.